



FIFTY POUNDS

FOLPET-SEVIN 4-4 D

Manufactured for Use by Professional Gardeners and Nurserymen as a Fungicide-Insecticide Combination for Certain Diseases and Insects on Ornamental Trees, Shrubs and Grapes.

(Contains *Phaltan (Folpet) and **Sevin)

ACTIVE INGREDIENTS:

*Folpet (N-(trichloromethylthio) phthalimide		 4.00%
**Carbaryl (1-Naphthyl	N-methylcarbamate)	 4.00%
INERT INGREDIENTS:		 92.00%
	•	100.00%

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

For best results, apply when air is calm, cover foliage with a light film of dust. Consult your local State Agricultural Extension Service for specific information as the timing, number and rate of applications will vary with local conditions. For control of certain diseases, apply as preventive measure when conditions are favorable and repeat at 7 to 10 day intervals as necessary. For insect control apply when insects are first noted and repeat at 7 to 10 day intervals as necessary. This product is suitable for application through ground dusters, airplane, or hand dusters.

Do not apply within 7 days following oil sprays. Do not apply during blocm when honey bees or other pollinators are active. Do not use on Boston lvy.

ROSES: For control of Black Spot, Powdery Mildew, Aphids, Boxelder Bug, Japanese Beetle, Lace Bugs, and Thrip. Apply 30 to 40 lbs. per acre or 34 to 1 lb. per 1,000 sq. ft. Make first application when young leaves first appear and repeat at 7 to 10 day intervals throughout the growing season.

ORNAMENTAL TREES AND SHRUBS: For control of leaf feeding insects, Japanese Beetle, Bagworms, Tent Caterpillars.

GRAPES: For control of black rot, downy mildew, powdery mildew, grape berry moth, grape leafhopper, grape root worm, and eight spotted forester, apply 75 lbs. per acre or 13/4 lb. per 1,000 sq. ft. To control black rot and downy mildew make an application just before bloom and repeat post-bloom applications as indicated below. Make first post-bloom application immediately after bloom and repeat (2nd

post-bloom application) 10-14 days after first post-bloom. For additional application repeat when needed usually the last week in July or first week in August. Generally three applications are necessary: post-bloom, 10-14 days later and mid-summer. No time limitation—may be applied up to harvest.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Store in cool place. Protect from excessive heat. Burn bag immediately when empty. Stay out of smoke. Do not contaminant feed or foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AGWAY INC.
FERTILIZER — CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201