

1 (of 2)
AGWAY PROMOTIONAL SERVICES

COPY

FOR: Maneb-Malathion 5.6-5D

DATE: 9/12/73

PGA—50 CHARACTERS
 ELITE—60 CHARACTERS

NET WEIGHT

AGWAY^R

Maneb-Malathion 5.6-5D

ACTIVE INGREDIENTS:

Maneb (Manganese ethylene bisdithiocarbamate)	5.6%
**Malathion	5.0%

INERT INGREDIENTS:

89.4%
<u>100.0%</u>

** O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION: (((18 pt. types))) Keep Out Of Reach Of
 Children (((12 pt. type)))

Harmful if swallowed. Avoid breathing of
 dust. Avoid contact with eyes, skin and clothing.
 Wash thoroughly after using. Avoid contamination of
 feed and foodstuffs.

This product is toxic to fish. Do not apply when weather
 conditions favor drift from areas treated. Keep out of
 any body of water. Do not contaminate water by cleaning
 of equipment or disposal of wastes. Do not reuse
 empty container. Destroy it by burying with waste
 or burning. Stay away from smoke or fumes.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF
 SALE. USE STRICTLY IN ACCORDANCE WITH LABEL
 CAUTION, WARNINGS AND DIRECTIONS.

For effective disease and insect control, make the
 first application when plants are small and repeat at

OK BY

SHEET NO 142

FOR: Maneb-Malathion 5.6-5D

DATE: 9/12/73

PCA—50 CHARACTERS
ELITE—60 CHARACTERS

7 to 10 day intervals as necessary. Do not exceed the dosage or apply beyond the date indicated under the crops listed. Use lower rate when plants are small.

CELERY: For control of Early Blight (*Cercospora*) and Late Blight (*Septoria*) and Aphids. Apply when plants are established in the field and repeat at 7 to 10 day intervals as long as blight threatens or insects are noted. Apply 20 to 25 lbs. per acre. Do not apply within 14 days of harvest. Remove excess residue by stripping, trimming and washing.

POTATOES: For control of Early and Late Blight, False Chinch Bug, Aphids and Leafhoppers. Start applications as soon as the plants are 2 to 6 inches high and repeat at 7 to 10 day intervals as long as blights threatens or insects are noted. Apply 20 to 40 lbs. per acre.

TOMATOES: For control of Early and Late Blight, *Septoria* Leaf Spot, Gray Leaf Spot (*Stemphylium*), Anthracnose, Aphids, Russet Mites, Flea Beetle and Leafhoppers. Apply soon after plants have been set to the field. Repeat applications at 7 to 10 day intervals as needed. Apply 20 to 35 lbs. per acre. Do not apply within 5 days of harvest.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

AGWAY INC., FERTILIZER-CHEMICAL DIVISION, BOX 1222 SYRACUSE NEW YORK 13201

EPA Reg. No.
8590-197

OF BY

SHEET NO 2/2