



8590-187



# COPPER-PARATHION 7-1.5 D

(A Fungicide, Insecticide Dust Combination for  
controlling certain Diseases and Insects on Potatoes)

## ACTIVE INGREDIENTS:

Copper (in Basic Copper Sulfate)  
expressed as metallic copper ..... 7.0  
Parathion (O, O-diethyl O-p-nitrophenyl thiophosphate) ..... 1.5%

INERT INGREDIENTS: ..... 91.5%  
-----  
100.0%



## POISON



## DANGER: KEEP OUT OF REACH OF CHILDREN.

May be fatal if swallowed, inhaled, or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes or on skin. Do not get on clothing. Wear natural rubber gloves, protective clothing, and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for parathion protection. Keep all unprotected persons out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated.

Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke or fumes.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

## DIRECTIONS

**READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.**

A fungicide, insecticide dust mixture for controlling certain diseases and insects affecting potatoes. This dust mixture contains Parathion which is not stable as a dust and should be used within one growing season.

**POTATOES:** For control of Early and Late Blight, Aphids, Blister Beetles, Leaf Hoppers and Thrips. Apply 30 to 40 lbs. per acre. Make the first application when conditions are favorable to the development of disease or when insects are first noted and repeat at 7 to 10 day intervals as necessary. Consult your State Agricultural Extension Service for recommendations under unusual conditions. Do not apply to potatoes within 5 days of harvest.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

Agway Inc., Chemical Division  
Box 1333, Syracuse, N. Y. 13201

EPA Reg. No. 8590-187

**NET CONTENTS FIFTY POUNDS**