


POLYRAM 7D®



NET CONTENTS 50 POUNDS

For Treating Seed Potatoes (Irish) only for the Control of Fusarium Seed Decay, Black Leg and Seed-Borne Scab.

ACTIVE INGREDIENTS:

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamate)] with 1 part by weight (16.1%) ethylenebis (thiocarbamic acid) bimolecular and trimolecular cyclic anhydrosulfides and disulfides 7.0%

INERT INGREDIENTS: 93.0%

TOTAL 100.0%

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin! In case of contact, flush with plenty of water; for eyes get medical attention. Do not breathe dust. Wear rubber gloves and respirator when mixing dust in hopper box. Do not use treated seed for feed or food.

This product is toxic to fish and wildlife. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Polyram 7D is effective as a Seed Piece Treatment for Irish Potatoes for control of Fusarium Seed decay, Black leg, and Seed-borne-scab. For best results, good coverage of seed pieces is necessary. Coat each seed piece evenly with a high film of dust immediately after cutting. If planting is delayed, store seed pieces in open crates two days before bagging.

POTATO SEED PIECE TREATMENT:

Rate: Use 1 to 1½ lbs. of dust to each 100 lbs. of cut seed pieces. Apply in convenient container by tumbling or by duster attachment over belt.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

Manufactured by Agway Inc., Syracuse, New York

USDA Reg. No. 8590-202
AC202F

®Registered Trademark of Badische Anilinund Soda-Fabrik AG. Ludwigshafen/Rhein