

15% EMULSIFIABLE,
KILLS MANY INSECTS
ON ORNAMENTALS,
FLOWERS AND
VEGETABLES

CAUTION:


KEEP OUT OF REACH
OF CHILDREN

See additional cautions and
warnings on back panel.

ACTIVE INGREDIENTS: Pct. by wt.
*Malathion 55.00%
Petroleum Distillate 35.00%
INERT INGREDIENTS,
not more than 10.00%
Total 100.00%

*O,O-dimethyl dithiophosphate of
diethyl mercaptosuccinate

MALATHION INSECT SPRAY



NET CONTENTS ONE QUART



CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and food-stuffs. Injury may occur on certain ferns such as Boston, Maidenhair and Pteris, as well as some species of Crassula. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. This product is toxic to fish.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

INSECTS CONTROLLED: Aphids, Asparagus beetles, Bagworms, Cabbage looper, Cottony scale, Forbes scale, Gall mites, Honey locust mites, Imported cabbage worms, Japanese beetle adults, Juniper scale, Lace bugs, Leafhoppers, Leaf miners, Lecanium scale, Magnolia scale, Mealy bugs, Mexican bean beetle adults, Oyster shell scale, Psyllids, Red mites, Scurfy scale, Spittlebugs, Spruce mites, Tarnished plant bugs, Tent caterpillars, Thrips, Two-spotted mites, Tulip tree scale.

APPLY TO FOLLOWING PLANTS AS SPECIFIED:
No limitations on applications to Flowers, Ornamentals and Potatoes.

On Asparagus, Beans, Cucumbers, Melons, Squash and Tomatoes, do not apply within 1 day of harvest.

On Broccoli, Eggplants, Onions, Peas, Peppers, Pumpkins, Strawberries and Turnips, do not apply within 3 days of harvest.

On Beets, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Kale, Head lettuce, Radishes, Spinach, do not apply within 7 days of harvest.

METHOD, RATE AND TIME OF APPLICATION:
Apply as a spray using 2 teaspoonsful of MALATHION INSECT SPRAY per gallon of water. Spray entire plant thoroughly, including lower leaf surfaces. Apply as insects appear. Repeat as needed.

OUTDOORS: Patios and similar areas for control of Mosquitoes and other small flying insects. Apply as spray using 1 tablespoon to a pint of water for 600 sq. feet of area. Spray Patios, Barbeque area, Building foundations, Shrubs, Trees and Grass Areas.

DOG AND CAT QUARTERS: For control of Fleas in Kennels, Pens, Yards, under Houses and Lawn areas. Use 5 ounces (10 table-spoonsful) to a gallon of water and apply at rate of 1 gallon of spray per 1000 sq. feet. Avoid contamination of food, feeders and water containers.

ANT MOUNDS: Use 1 tablespoon to a pint of water. Spray Ant hills thoroughly and mix moist earth in the top of the hill. For other small ants in Flower beds, Lawns and around Trees, spray lightly in the infested area. Repeat in 10 to 15 days if ants return.

NOTE: This pesticide is to be sold ONLY in this original unopened package.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk for any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

MANUFACTURED BY AGWAY INC.
CHEMICAL DIVISION,
BOX 1333, SYRACUSE, N. Y. 13201
USDA Reg. No. 8590-228

8590-228

01/22/69

1/1