

8590-228

02/21/70

174

AGWAY PROMOTIONAL SERVICES

COPY

FOR: LABEL COPY - MALATHION INSECT SPRAY

DATE: 12/4/69

((FRONT PANEL))

((AGWAY T.M.))

NET CONTENTS

MALATHION INSECT SPRAY

55% Emulsifiable, Kills Many Insects  
on Ornamentals, Flowers and Vegetables

ACTIVE INGREDIENTS: Percent by Wt.

*Malathion . . . . .	55.00%
Xylene . . . . .	35.00%

INERT INGREDIENTS, not more than . . . 10.00%

TOTAL 100.00%

\*O,O-dimethyl dithiophosphate of diethyl mercapto-  
succinate (Premium grade)

CAUTION: ((18 pt. type)) KEEP OUT OF REACH OF  
CHILDREN. ((12 pt. type)) See additional cautions  
and warnings on ((fill in)) panel.

**ACCEPTED**

FEB 21 1970

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND ROENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 8590-228 SUBJECT  
NO ATTACHED COMMENTS.

O.K BY lt

FOR: LABEL COPY - MALATHION INSECT SPRAY

DATE: 12/4/69

(((REAR PANEL)))

CAUTION: Harmful if swallowed. Avoid prolonged breathing of spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

Injury may occur on certain ferns such as Boston, Maidenhair and Pteris, as well as some species of Crassula.

This product is toxic to fish. Keep out of any body of water. Do not reuse container. Destroy it by burying or burning (keep away from smoke).

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

INSECTS CONTROLLED: Aphids, Asparagus beetles, Bagworms, Cabbage looper, Cottony scale, Forbes scale, Gall mites, Honey locust mites, Imported cabbage worms, Japanese beetle adults, Juniper scale, Lace bugs, Leafhoppers, Leaf miners, Lecanium scale, Magnolia scale, Mealy bugs, Mexican bean beetle adults, Oyster shell scale, Psyllids, Red mites, Scurfy scale, Spittlebugs, Spruce mites, Tarnished plant bugs, Tent caterpillars, Thrips, Two-spotted mites, Tulip tree scale.

## APPLY TO FOLLOWING PLANTS AS SPECIFIED:

No limitations on applications to Flowers, Ornamentals and Potatoes.

On Asparagus, Beans, Cucumbers, Melons, Squash and Tomatoes, do not apply within 1 day of harvest.

On Broccoli, Eggplants, Onions, Peas, Peppers, Pumpkins, Strawberries and Turnips, do not apply within 3 days of harvest.

On Beets, Brussels Sprouts, Cabbage, Carrots, Cauliflower, Kale, Head lettuce, Radishes, Spinach, do not apply within 7 days of harvest.

OK BY lt

FOR: LABEL COPY - MALATHION INSECT SPRAY

DATE: 12/4/68

((( REAR PANEL CONTINUED )))

METHOD, RATE AND TIME OF APPLICATIONS: Apply as a spray using 2 teaspoonsful of MALATHION INSECT SPRAY per gallon of water. Spray entire plant thoroughly, including lower leaf surfaces. Apply as insects appear. Repeat as needed.

OUTDOORS: Patios and similar areas for control of Mosquitoes and other small flying insects. Apply as spray using 1 tablespoon to a pint of water for 600 sq. ft. of area. Spray Patios, Barbeque area, Building foundations, Shrubs, Trees and Grass Areas.

DOG AND CAT QUARTERS: For control of Fleas in Kennels, Pens, Yards, under Houses and Lawn areas. Use 5 ounces (10 tablespoonsful) to a gallon of water and apply at rate of 1 gallon of spray per 1000 sq. ft. Avoid contamination of food, feeders and water containers.

ANT MOUNDS: Use 1 tablespoon to a pint of water. Spray Ant hills thoroughly and mix moist earth in the top of the hill. For other small ants in Flower beds, Lawns and around Trees, spray lightly in the infested area. Repeat in 10 to 15 days if ants return.

STORED GRAIN: In barley, corn, grain sorghum, oats, rye, wheat and garden seeds for control of confused flour beetle, granary weevil, saw-toothed grain beetle, flat grain beetle, red flour beetle, rusty grain beetle, lesser grain borer and Indian meal moth apply Malathion Insect Spray as follows.

Residual spray for treating surfaces prior to storage:  
Apply to walls, floors, machinery in grain elevators, truck beds, box cars or ships' holds using 1 gallon of Malathion Insect Spray in 25 gallons of water. Make a complete coverage application after thoroughly cleaning the area to be sprayed. Remove and burn all sweepings and debris.

REGISTERED CHARACTER  
UNREGISTERED CHARACTER

O.K. BY It

4/4

AGWAY PROMOTIONAL SERVICES

FOR: LABEL COPY - MALATHION INSECT SPRAY

DATE: 12/4/69

(((REAR PANEL CONTINUED)))

Grains going into storage: Apply 1 pint of Malathion Insect Spray in 2-3 gallons of water per 1,000 bushels. Apply as the grain is being loaded or being turned into final storage.

After grains are stored: To control Indian meal moth apply 1/2 pint of Malathion Insect Spray in 1-2 gallons of water per 1,000 square feet of grain surface area. Apply evenly over the grain surface immediately after storing and repeat if necessary.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk for any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

MANUFACTURED BY AGWAY INC., CHEMICAL DIVISION,  
BOX 1333, SYRACUSE, N.Y. 13201

USDA Reg. No. 8590-228

OK BY It