


UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR EFF. SUBJECT
 ED UNDER NO 8590-22
 TO ATTACHED COMMENTS.

**KILLS MANY INSECTS
 ON ORNAMENTALS,
 FLOWERS AND
 VEGETABLES**

CAUTION:
 KEEP OUT OF REACH
 OF CHILDREN
 See additional cautions and
 warnings on back panel.

ACTIVE INGREDIENTS: Pct. by wt.
 *Malathion 55.00%
 Xylene 35.00%
INERT INGREDIENTS,
 not more than 10.00%
 Total 100.00%
 *O, O-dimethyl dithiophosphate of
 diethyl mercaptosuccinate (Premium
 grade)

**MALATHION INSECT
 SPRAY**



CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin; wash thoroughly using. Avoid contamination of feed and foodstuffs. Injury may occur on certain ferns such as Boston fern and Pteris, as well as some species of Crassulaceae. This product is toxic to fish. Keep out of any water. Do not reuse container. Destroy it by burning (keep away from smoke). This product is highly toxic to bees exposed to treatment or residues on crops. Protective clothing may be obtained from your cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL OR STORE NEAR OPEN FLAME.

DIRECTIONS
 READ THESE ENTIRE DIRECTIONS AND WARNINGS. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS. ONLY THE FOLLOWING CROPS AS SPECIFIED. Limitations on applications to Flowers, Ornamentals, Potatoes.

ASPARAGUS --- Controls asparagus beetle and spider mites. Do not apply within 1 day of harvest. **BEANS** --- Controls Mexican bean beetle, leafhoppers, spider mites, beetle, aphids and cucumber beetle. Do not apply within 1 day of harvest. **COLE CROPS & LEAFY VEG** (BROCCOLI, BRUSSELS SPROUTS, CABBAGE, COLLARDS, KALE, TURNIPS) --- Controls aphids, caterpillars (on cauliflower, collard, brussels sprouts and only), and Harelequin cabbage bugs (on collard). Do not apply to broccoli within 3 days of harvest. Do not apply to brussels sprouts, cabbage, collards, kale or cauliflower within 7 days of harvest. **CAULIFLOWERS** --- Controls aphids and caterpillars. Do not apply within 7 days of harvest. **CUCUMBERS** --- Controls aphids and spider mites. Do not apply within 1 day of harvest. **EGGPLANTS** --- Controls aphids, and spider mites. Do not apply within 30 days of harvest. **HEAD LETTUCE** --- Controls aphids, spider mites and leafhoppers. Do not apply within 7 days of harvest. **MELONS** (CANTALOUPE, CASABABA, SHAWNS, HONEY DEWS, MUSKMELONS, AND MELONS) --- Controls aphids, spider mites and beetles. Do not apply within 1 day of harvest. **ONIONS** --- Controls thrips and onion maggot. Do not apply within 3 days of harvest. **PEPPERS** --- Controls aphids and spider mites. Do not apply within 3 days of harvest. **POTATOES** --- Controls aphids, and mites. Do not apply within 1 day of harvest. **ROOT CROPS** (BEETROOTS, CARROTS, RUTABAGAS) --- Controls aphids. Do not apply within 1 day of harvest. **SQUASH** --- Controls aphids, spider mites and pickleworm. Do not apply within 1 day of harvest.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

Injury may occur on certain ferns such as Boston, Maiden-hair and Pteris, as well as some species of Crassula.

This product is toxic to fish. Keep out of any body of water. Do not reuse container. Destroy it by burying or burning (keep away from smoke).

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

APPLY TO FOLLOWING PLANTS AS SPECIFIED. No limitations on applications to Flowers, Ornamentals and Potatoes.

ASPARAGUS — Controls asparagus beetle and thrips.

Do not apply within 1 day of harvest. **BEANS** — Controls mexican bean beetle; leafhoppers, spider mites, Japanese beetle, aphids and cucumber beetle. Do not apply within 1 day of harvest.

COLE CROPS & LEAFY VEGETABLES (BROCCOLI, BRUSSELS SPROUTS, CABBAGE, COLLARDS, KALE, TURNIPS) — Controls aphids, caterpillars (on cauliflower collard, brussels sprouts and broccoli only), and Harelequin cabbage bugs (on collards only). Do not apply to broccoli within 3 days of harvest. Do not apply to brussels sprouts, cabbage, collards, kale, turnips, cauliflower within 7 days of harvest.

CAULIFLOWER — Controls aphids and caterpillars. Do not apply within 7 days of harvest. **CUCUMBERS** — Controls aphids, spider mites. Do not apply within 1 day of harvest.

EGGPLANTS — Controls aphids, and spider mites. Do not apply within 30 days of harvest. **HEAD LETTUCE** — Controls aphids, mites and leafhoppers. Do not apply within 7 days of harvest.

MELONS (CANTALOUPS, CASABAS, CRENSHAW, HONEY DEWS, MUSKMELONS, AND WATER MELONS) — Controls aphids, spider mites and cucumber beetles. Do not apply within 1 day of harvest.

ONIONS — Controls thrips and onion maggots. Do not apply within 3 days of harvest. **PEPPERS** — Controls aphids and pepper maggot. Do not apply within 3 days of harvest.

PUMPKINS — Controls aphids, and mites. Do not apply within 3 days of harvest. **ROOT CROPS (BEETS, CARROTS, AND RADISHES)** — Controls aphids. Do not apply within 7 days of harvest.

SQUASH — Controls aphids, spider mites and pickleworm. Do not apply within 1 day of harvest. **STRAW-**

BERRIES — Controls aphids, spider mites, Lygus bug, spittlebugs, field crickets and thrips. Do not apply within 3 days of harvest. To control strawberry root weevil apply 2-4 teaspoons per gallon of water. Apply this to the soil surface before planting and work into the top 6 to 8 inches of soil.

TOMATOES (OUTDOOR) — Controls spider mites, aphids, and Drosophila. Do not apply within 1 day of harvest. **SPINACH** — Controls aphids. Do not apply within 7 days of harvest.

ORNAMENTALS & FLOWERS — Controls Oyster shell scale, lace bug, Euonymus scale, aphids, mealybugs, spider mites, whitefly, four-lined leaf bug, Japanese beetle adults, potato leafhopper, tarnished plant bug, thrips, rose leafhopper, European pine shoot moth, scurfy scale, birch, leaf miner, boxwood leaf miner, bagworms, tent caterpillar, Azalea scale, oak kermes*, pine leaf scale, magnolia scale, Fletcher scale*, Florida red scale*, juniper scale*, black scale crawlers, wax scale.

*Apply when scale crawlers have settled on foliage.

METHOD RATE, AND TIME OF APPLICATION: Apply as a spray using 2 teaspoonful of MALATHION INSECT SPRAY per gallon of water. Spray entire plant thoroughly, including lower leaf surfaces. Apply as insects appear. Repeat as needed.

CURTAINS: Patios and similar areas for control of Mosquitoes and other small flying insects. Apply as spray using 1 tablespoon to a pint of water for 600 sq. feet of area. Spray Patios, Barbeque area, Building foundations, Shrubs, Trees and Grass Areas.

DOG AND CAT FLIES: For control of Fleas in Kennels, Pens, Yards, under Houses and Lawn areas. Use 5 ounces (10 tablespoonsful) to a gallon of water and apply at rate of 1 gallon of spray per 1000 sq. feet. Avoid contamination of food, feeders and water containers.

ANT HILLS: Use 1 tablespoon to a pint of water. Spray Ant hills thoroughly and mix moist earth in the top of the hill. For other small ants in Flower beds, Lawns and around trees, spray lightly in the infested area. Repeat in 10 to 15 days if ants return.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk for any such use. Any damage arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

MANUFACTURED BY AGWAY INC. CHEMICAL DIVISION
BOX 1333, SYRACUSE, N.Y. 13201 USDA Reg. No. 8590-228