

Reg # 8590-228

PM-14

197

AGWAY
MALATHION INSECT SPRAY

Kills aphids, mealybugs, scale insects and mosquitoes on vegetables, flowers and ornamentals.

| ACTIVE INGREDIENT: | % by Weight |
|---|---------------|
| Malathion: 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate | 55.00% |
| INERT INGREDIENTS* | <u>45.00%</u> |
| TOTAL | 100.00% |

*Contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements on back panel.

Net Contents

Agway, Inc.
P.O. Box 4741
Syracuse, NY 13221


EPA Reg. No. 8590-228

EPA Est. No. 1386-OH-1

red. P

to be

to



287

ADVERTISING AND SALES PROMOTION SERVICES

33PV

FOR: INSECT SPRAY WITH MALATHION

DATE: 10/27/88

((BACK PANEL))

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin. Wash thoroughly after using. Avoid contamination of food and feed.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water. See a physician if irritation persists.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL CHEMICAL HAZARDS

Do not use or store near heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a

OK BY
.....
.....

487

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: Insect Spray with Malathion

DATE: 10/27/88

APPLY TO FOLLOWING PLANTS AS SPECIFIED:

PICA - 50 CHARACTERS
ELITE - 50 CHARACTERS

VEGETABLES

ASPARAGUS - Controls asparagus beetle and thrips.
Do not apply within 1 day of harvest.

BEANS - Controls mexican bean beetle, leafhoppers,
spider mites, Japanese beetle, aphids and cucumber
beetle. Do not apply within 1 day of harvest.

COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CABBAGE,
COLLARDS, KALE, TURNIPS) - Controls aphids, cater-
pillars (on cauliflower, collard, brussels sprouts
and broccoli only), and Harelequin cabbage bug
(on collards only). Do not apply to broccoli
within 3 days of harvest. Do not apply to brussels
sprouts, cabbage, collards, kale, turnips, cauli-
flower within 7 days of harvest.

CAULIFLOWER - Controls aphids and caterpillars.
Do not apply within 7 days of harvest.

CUCUMBERS - Controls aphids, spider mites. Do not
apply within 1 day of harvest.

OK BY
.....
.....

EGGPLANTS - Controls aphids, and spider mites. Do
not apply within 3 days of harvest.

LETTUCE - Controls aphids, mites and leafhoppers.
Do not apply within 7 days of harvest on head

597

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: Insect Spray with Malathion

DATE: 10/27/88

MELONS (CANTALOUPE, CASABAS, CRENSHAW, HONEY DEWS, MUSKMELONS, AND WATERMELONS) - Controls aphids, spider mites and cucumber beetles. Do not apply with 1 day of harvest. Do not apply for cucumber beetles unless plants are dry.

ONIONS - Controls thrips and onion maggots. Do not apply with 3 days of harvest.

PEPPERS - Controls aphids and pepper maggot. Do not apply within 3 days of harvest.

PUMPKINS - Controls aphids and mites. Do not apply within 3 days of harvest.

ROOT CROPS (BEETS, CARPOTS, AND RADISHES) - Controls aphids. Do not apply within 7 days of harvest.

SQUASH - Controls aphids, spider mites and pickle worm. Do not apply within 1 day of harvest.

STRAWBERRIES - Controls aphids, spider mites, strawberry root weevil, Lygus bug, spittlebugs, field crickets and thrips. Do not apply within 3 days of harvest.

TOMATOES (OUTDOOR) - Controls spider mites, aphids

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

COPY
.....
.....

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR: Insect Spray with Malathion

DATE: 10/27/88

SPINACH - Controls aphids. Do not apply within 7 days of harvest.

PICA 50 CHARACTERS ELITE - 60 CHARACTERS

ORNAMENTALS & FLOWERS - Controls Oyster shell scale, lace bug, Euonymus scale, aphids, mealybugs, spider mites, whitefly, four-lined leaf bug, Japanese beetle adults, potato leafhopper, tarnished plant bug, thrips, rose leafhopper, European pine shoot moth, scurfy scale, birch leaf miner, boxwood leaf miner, bagworms, tent caterpillar, Azalea scale, oak kermes*, pine leaf scale, magnolia scale, Fletcher scale*, Florida red scale*, juniper scale*, black scale crawlers, wax scale.

*Apply when scale crawlers have settled on foliage.

Injury may occur on certain ferns such as Boston, Maidenhair and Pteris, as well as some species of Crassula, Canaerti Juniper, Viburnum lantana, Hickory, Small-leaf Spirea, White Pine and Maples.

OUTDOORS: Patios and similar areas for control of mosquitoes and other small flying insects.

OK BY
.....
.....

Apply spray using 1 tablespoon to a pint of water per 100 sq. ft. of area. Spray patios, barbecue area, building foundations, shrubs, trees and grass areas.

DOG AND CAT QUARTERS: For control of fleas in kennels, pens, yards, under houses and lawn area.

7 37

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: Insect Spray with Malathion

DATE: 10/27/88

water and apply at rate of 1 gallon of spray per 1000 sq. ft. Avoid contamination of food, feeders and water containers.

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

ANT MOUNDS: Use 1 tablespoon to a pint of water. Spray Ant hills thoroughly and mix moist earth in the top of the hill. For other small ants in flower beds, lawns and around trees, spray lightly in the infested area. Repeat in 10 to 15 days if ants return.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

OK BY

•••••
•••••