

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

FLOWERS AND EVERGREENS: This material may be applied as a dust or spray for the control of such diseases as powdery mildew on roses, phlox, zinnias, sweet peas and lilacs; rust on snapdragons, hollyhock and carnations; and to aid in keeping red spiders in check on evergreens. For red spiders delay application until the new growth has hardened. Apply every week or ten days during the growing season, or more often during rainy weather. Cover all parts of the plant with a thin film of dust or spray.

PEACHES, CHERRIES, PLUMS AND PRUNES: On stone fruits for brown rot, apply just as trees are coming into bloom, and after the petals fall, and at intervals of two weeks during season. During the last month before harvest, dust lightly every week or ten days.

APPLES AND PEARS: For scab on apples or pears dust or spray weekly from the time the first green tissue shows until 30 days after petals fall. Use more frequently during rainy periods.

DUSTING: This material is ready for use and should be applied with a duster. Cover plants lightly from two sides when the air is quiet.

SPRAYING: Mix 4 tablespoonsful in each gallon of water and cover all leaf surfaces with a fine mist. With garden sprayers, shake frequently for agitation. In large sprayers mix 5 to 6 pounds in 100 gallons of water.

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

FLOWERS AND EVERGREENS: This material may be applied as a dust or spray for the control of such diseases as powdery mildew on roses, phlox, zinnias, sweet peas and lilacs; rust on snapdragons, hollyhock and carnations; and to aid in keeping red spiders in check on evergreens. For red spiders delay application until the new growth has hardened. Apply every week or ten days during the growing season, or more often during rainy weather. Cover all parts of the plant with a thin film of dust or spray.

PEACHES, CHERRIES, PLUMS AND PRUNES: On stone fruits for brown rot, apply just as trees are coming into bloom, and after the petals fall, and at intervals of two weeks during season. During the last month before harvest, dust lightly every week or ten days.

APPLES AND PEARS: For scab on apples or pears dust or spray weekly from the time the first green tissue shows until 30 days after petals fall. Use more frequently during rainy periods.

DUSTING: This material is ready for use and should be applied with a duster. Cover plants lightly from two sides when the air is quiet.

SPRAYING: Mix 4 tablespoonsful in each gallon of water and cover all leaf surfaces with a fine mist. With garden sprayers, shake frequently for agitation. In large sprayers mix 5 to 6 pounds in 100 gallons of water.

AGWAY

wettable sulfur dust or spray

A SULFUR FUNGICIDE FOR USE
ON FRUITS AND FLOWERS

ACTIVE INGREDIENTS:	Percent by Wt.
Sulfur, Average particle size, 3 to 5 microns	95.00%
INERT INGREDIENTS, not more than	5.00%
Total	100.00%

caution:

KEEP OUT OF REACH OF CHILDREN

See additional cautions and warnings on side panel.

NET WEIGHT 4 LBS.

ACCEPTED

FEB 5 1971

UNDER FEDERAL INSECTICIDE
AND RODENTICIDE ACT
FOR FEDERAL POISON REGISTER-
ED UNITS 8597-264

CAUTION: Do not use sulfur on apple foliage within two weeks of an oil spray. Avoid dusting or spraying when temperature is above 85°F.

Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.