



scale-rid

(DORMANT SPRAY OIL WITH ETHION*)

(COMBINES ETHION WITH SUPERIOR OIL FOR CONTROL OF CERTAIN INSECTS ON ORNAMENTAL TREES AND SHRUBS AND FRUIT TREES)

ACTIVE INGREDIENTS:

*Ethion 2.0%

**Petroleum Oil (Superior Type 60 Second Oil) 95.0%

INERT INGREDIENTS 3.0%

100.0%

*O,O,O',O'-Tetraethyl S,S'-methylene bisphosphorodithioate U.S. Pat. No. 2,873,228

THIS PRODUCT CONTAINS—.15 lb. Ethion Per Gallon

****Superior Oil Specifications:**

Unsulfonated Residue 92-94%

Viscosity 60 seconds

DO NOT STORE AT TEMPERATURES BELOW 10°F.

caution:

KEEP OUT OF REACH OF CHILDREN

See additional cautions and warnings on back panel.

NET CONTENTS

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products. Do not apply to any plant during bloom period. May be injurious for resident birds. This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from crop areas treated.

NOTE: Do not use Scale Rid on oil sensitive plants such as Sugar and Japanese Maple, Beech, Birch, Hickory, Walnut and Butternut. Oil will injure Douglas Fir flower buds and remove temporarily blue color from Colorado Blue Spruce.

general information

Scale Rid combines the spreading and penetrating advantages of oil with the long residual insecticide Ethion which is effective for use against Scale insects, Mites and Aphids. Scale Rid can be used as a Dormant or Delayed Dormant Spray.

Apply Scale Rid during Dormant or Delayed Dormant Period. Thorough coverage is necessary for good control. Fill the sprayer with water, add the desired amount of Scale Rid, shake, stir or agitate until thoroughly mixed. Do not apply Captan, Folpet (Phaltan), or Sulphur within 10 days of any spray containing oil.

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

DORMANT AND DELAYED DORMANT SPRAYS

Apply early spring when temperature is not likely to fall below freezing before the spray has had a chance to dry. The temperatures at time of application should be above 40 F.

DEFINITIONS: DORMANT SPRAY: Apply before buds swell **SILVER-TIP SPRAY:** Apply when leaves of blossom buds are breathing **DELAYED DORMANT SPRAY:** Apply when leaves of the blossom buds are out 1/4 to 1/2.

APPLES AND PEARS: For control of European Red Mite, Apple Aphid, Rosy Apple Aphid, San Jose and Oyster Shell Scale. Apply as a Dormant or Delayed Dormant Spray (see definitions above) for over wintering EGGS of mites and aphids. Mix 5 level tablespoonsful per gallon of water, or 1 quart per 12 1/2 gals. of water, or 2 gals. per 100 gals. of water. Spray thoroughly to the point of run off.

CHERRIES: For control of San Jose Scale and Black Cherry Aphid apply as a Dormant Spray. Mix 5 level tablespoonful per gallon of water.

NECTARINES, PEACHES, PLUMS AND PRUNES: For control of San Jose Oyster Shell Scale, Brown Almond and European Red Mites and Aphid apply as Silver Tip Spray for control of scale insects and over wintering Mite and Aphid Eggs. Mix 5 level tablespoonful per gallon of water, or 1 quart per 12 1/2 gallons of water, or 2 gallons per 100 gallons of water. Spray thoroughly to the point of run off.

Apr. 6, 1970
 234

OR
 Ma
 Will
 For
 Tra
 Sca
 2-3
 For
 Scu
 locl
 or
 Ap
 nig
 No
 aft
 eas
 Be
 juri
 Col
 PIN
 tro
 ful
 wal
 hat
 ger
 Au
 Apl
 lon
 rial
 Blu
 AZI
 TO
 For
 red
 cra
 da
 or
 ag
 WA
 ch
 sta
 um
 noi
 pal
 of
 noi
 see
 us
 sh
 col
 val
 NO
 un
 MA
 CH
 US

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid storage near feed or food products. Do not apply to any plant during bloom period. May be injurious for resident birds.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label. Do not apply when weather conditions favor drift of spray from crop areas treated.

NOTE: Do not use Scale Rid on oil sensitive plants such as Sugar and Japanese Maple, Beech, Birch, Hickory, Walnut and Butternut. Oil will injure Douglas Fir flower buds and remove temporarily blue color from Colorado Blue Spruce.

general information

Scale-Rid combines the spreading and penetrating advantages of oil with the long residual insecticide Ethion which is effective for use against Scale insects, Mites and Aphids. Scale-Rid can be used as a Dormant or Delayed Dormant Spray.

Apply Scale-Rid during Dormant or Delayed Dormant Period. Thorough coverage is necessary for good control. Fill the sprayer with water, add the desired amount of Scale-Rid, shake, stir or agitate until thoroughly mixed. Do not apply Captan, Folpet (Phaltan), or Sulphur within 10 days of any spray containing oil.

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

DORMANT AND DELAYED DORMANT SPRAYS

Apply early spring when temperature is not likely to fall below freezing before the spray has had a chance to dry. The temperatures at time of application should be above 40°F.

DEFINITIONS: DORMANT SPRAY: Apply before buds swell. **SILVER-TIP SPRAY:** Apply when leaves of blossom buds are breathing. **DELAYED DORMANT SPRAY:** Apply when leaves of the blossom buds are out 1/4 to 1/2".

APPLES AND PEARS: For control of European Red Mite, Apple Aphid, Rosy Apple Aphid, San Jose and Oyster-Shell Scale. Apply as a Dormant or Delayed Dormant Spray (see definitions above) for over wintering eggs of mites and aphids. Mix 5 level tablespoonsful per gallon of water, or 1 quart per 12 1/2 gals. of water, or 2 gals. per 100 gals. of water. Spray thoroughly to the point of run-off.

CHERRIES: For control of San Jose Scale and Black Cherry Aphid apply as a Dormant Spray. Mix 5 level tablespoonsful per gallon of water.

NECTARINES, PEACHES, PLUMS AND PRUNES: For control of San Jose, Oyster-Shell Scale, Brown Almond and European Red Mites and Aphid, apply as Silver-Tip Spray for control of scale insects and over-wintering Mite and Aphid Eggs. Mix 5 level tablespoonsful per gallon of water, or 1 quart per 12 1/2 gallons of water, or 2 gallons per 100 gallons of water. Spray thoroughly to the point of run-off.

ORNAMENTALS: For control of Scale insects on Maple, Elm, Catalpa, Flowering Quince, Ash, Red Willow, Dogwood, etc. For control of Mites on Hemlock.

For Control of Soft Scale on Maple, Elm, Tree Cottony Maple, etc. Mix 7 level tablespoonsful per gallon of water or 2 3/4 gallons of Scale-Rid per 100 gallons of water.

For Control of Hard or Armored Scales, Oyster-Shell Scurfy, San Jose, Pine-needle, Juniper, Euonymus, Hemlock Scale. Mix 5 level tablespoonsful per gallon of water or 2 gallons of Scale-Rid per 100 gallons of water.

Apply as dormant spray after most danger of freezing nights has passed but before buds have begun to open. Note: Spray on clear mid day and do not spray in late afternoon. Be cautious with oil sprays. Do not use on easily injured plants such as Sugar and Japanese Maple, Beech, Birch, Hickory, Walnut and Butternut. Oil will injure Douglas Fir flower buds and remove blue color from Colorado Blue Spruce.

PINES, SPRUCE, FIR, CEDAR AND HEMLOCK: For control of Hemlock scale. Apply 1/4 pint (1 1/2 standard cupsful) per 25 gallons or 1 1/2 teaspoonsful per gallon of water. Apply mid-spring when overwintering eggs start to hatch. Repeat in 10 to 14 days if necessary. A second generation occurs mid August. Repeat applications in August if necessary. For control of Eastern Spruce Gall Aphid. Apply early April at rate of 1 1/2 pints per 25 gallons of water or 1 1/2 teaspoonsful per gallon. This material will temporarily remove the blue color from Colorado Blue Spruce.

AZALEAS, CAMELLIAS, GARDENIAS, LIGUSTRUM PIT-TOSPORUM, SASANQUA, VIBURNUM, BOXWOOD, YEW: For control of Camellia scale, Euonymus scale, Florida red scale, Florida wax scale, Spider Mite. Apply when crawlers or Mites are first noted and repeat in 10 to 14 days if necessary. Apply 3 qts. per 100 gallons of water or 1 1/2 teaspoonsful per gallon of water. Thorough coverage is essential to both upper and lower leaf surfaces.

WARRANTY: Seller warrants that product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits, values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

MANUFACTURED BY AGWAY, INC.
CHEMICAL DIVISION, BOX 1333, SYRACUSE, N.Y. 13201
USDA Reg. No. 8590-234

8590-234

