

Agway Insect Dust with Sevin

For control of insects on vegetables,
fruits, ornamentals, and lawns



ACCEPTED
OCT 12 1977

KEEP OUT OF
REACH OF CHILDREN

CAUTION:

See other cautions on back panel

Net Wt. 1 lb.
(.45 kg)

| | |
|--|--------|
| ACTIVE INGREDIENTS: | |
| Carbaryl (1-naphthyl N-methylcarbamate) | 5.0% |
| INERT INGREDIENTS: | 95.0% |
| | 100.0% |

*Trademark of Union Carbide

CAUTION: Harmful if swallowed or inhaled. Avoid prolonged or repeated breathing of dust. Do not take internally. Skin contact with this insecticide may be harmful. Avoid contact. Wash hands and face before eating. Avoid contamination of food. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

General Information

Apply Agway Insect Dust when insects first appear, preferably before damage from feeding occurs. Repeat at 7 to 10 day intervals as necessary. Apply with conventional dusting equipment. Thorough coverage to top and underside of leaves is necessary. Avoid using during flowering and pollination period. Sevin is toxic to honey bees. Do not dust too heavily or crop injury may result to tender foliage in the presence of rain or high humidity following dusting.

Agway Insect Dust does not control mites. Where mites are a problem, use a suitable miticide.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Vegetables

ASPARAGUS: For control of asparagus beetle on seedlings and spears, apply ½ to ¾ lbs. per 1,000 sq. ft. when insects appear and repeat as needed, but not more often than once every three days. Do not apply within 1 day of harvest.

BEANS: For control of Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Western Bean Cutworm, Flea Beetles and Japanese Beetles: Apply ½ lb. per 1,000 sq. ft. For control of Corn Earworm, Armyworms and Velvet Bean Caterpillar: Apply ½ to ¾ lb. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 1 lb. per 1,000 sq. ft. May be applied up to and including the day of harvest on Beans.

CORN: For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle and Japanese Beetle. Apply ½ to 1 lbs. per 1,000 sq. ft. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to day of harvest on Corn.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI: For control of Flea Beetle and Harlequin Bug, Corn Earworm, Armyworm and Cabbage Caterpillars, apply ½ to ¾ lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest. Repeat at 7 to 10 day intervals as needed.

CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE (ESCAROLE), GARDEN BEETS (TOPS), HORSEADISH, KALE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY (TOPS), SPINACH, SWISS CHARD AND TURNIPS (TOPS): For control of Flea Beetles, Harlequin Bug and Leafhoppers apply ½ to ¾ lb. per 1,000 sq. ft. when insects first appear and repeat at 5 to 7 day intervals as needed. Do not apply within 14 days of harvest when leaves or tops are used. Where only roots of Beets, Radish, Horseradish, Parsnips, Rutabagas, Salsify, and Turnips are used, do not apply within 3 days of harvest.

CUCUMBERS, SQUASH (SUMMER AND WINTER), PUMPKINS, MELONS: For control of Cucumber Beetle, Melonworm, Pickleworm, Squash Bugs, and Flea Beetle, apply ½ lb. per 1,000 sq. ft. Apply when insect appears and repeat at 10 day intervals as needed. Avoid excessive applications. Some leaf injury may occur if treatment is made when tender foliage is wet or in presence of very high humidity.

LETTUCE: For control of Armyworm, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus and Stink Bugs: Apply ¾ lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals. Do not apply within 3 days of harvest on Head Lettuce or 14 days of harvest on Leaf Lettuce.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS: For control of Colorado Potato Beetle, Flea Beetle and Leafhoppers: Apply ½ to ¾ lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs and Stink Bugs: May be applied up to and including day of harvest. Apply 1 lb. per 1,000 sq. ft.

Fruits

BLUEBERRIES: For control of Japanese Beetle, Cherry and Cranberry Fruitworms, apply dust thoroughly covering top and underside of leaves. Apply ¾ lb. per 1,000 sq. ft. when insects are first noticed and repeat at 7 to 10 day intervals as necessary. May be applied up to and including the day of harvest.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, AND RASPBERRIES: For control of Japanese Beetle, Leaf Rollers, Leafhoppers and Raspberry Aphid, dust thoroughly covering top and underside of the leaves. Apply ¾ lb. per 1,000 sq. ft. when insects appear and repeat at 7 to 10 day intervals as needed. Full dust coverage is essential for good control. Do not apply within 7 days of harvest.

GRAPES: For control of Grape Berry Moth, Japanese Beetle, Grape Leaf Folder and Leafhoppers, dust thoroughly covering top and underside of leaves. Apply ¾ to 1 lb. per 1,000 sq. ft. when insects appear and repeat at 10 day intervals as needed. May be applied up to and including the day of harvest.

STRAWBERRIES: For control of Meadow Spittle Bug and Strawberry Leaf Roller, dust thoroughly covering top and underside of leaves. Apply ¾ lb. per 1,000 sq. ft. when insects first appear and repeat at 10 day intervals as needed. Do not apply within one day of harvest.

Ornamentals

For control of Aphids, Bagworm, Blister Beetle, Boxelder Bug, Elm Leaf Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Periodical Cicada, Plant Bugs, Scale Insect Crawlers, Tent Caterpillars, Exposed Thrips, dust thoroughly both top and underside of leaves. Apply at 7 to 10 day intervals throughout entire growing season, starting before serious build-up of insects occurs. Sevin has been used at recommended rates without damage to Andromeda, Arborvitae, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Camellia, Carnation, Chinese Elm, Chrysanthemum, Dahlia, Delphinium, Dogwood, Elm, Gladiolus, Hemlock, Holly, Hydrangeas, Iris, Juniper, Laurel, Lilac, Magnolia, Maidenhair Fern, Maple, Marigold, Nasturtium, Oak, Ornamental Mulberry, Petunia, Privet, Rhododendron, Rose, Rose of Sharon, Spirea, Verbena, Viburnum, Willow, Wisteria, Zinnia. Apply 1 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do NOT use on Boston Ivy.

Lawns

For control of Chinch Bugs, Fall Armyworms, Leafhoppers, Sod Webworms or Lawn Moths, Ants, Earwigs, Fleas, Millipedes, and Mosquitoes. Apply evenly 5 lbs. dust per 1,250 sq. ft. or 1 lb. per 250 sq. ft. For control of Chinch Bugs, Fall Armyworms, Sod Webworms and Ants apply water after application to move dust into the soil where insects frequent. Repeat in 7 to 14 days if necessary.

DISCLAIMER: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.



Manufactured by
Agway Inc.
Fertilizer-Chemical Division
Box 4933, Syracuse NY 13221

EPA Reg. No. 8590-257 EPA Est. No. 8590-PA-1

8590-257

10-12-1977

1/1