

8590-257 1061 PM12

AGWAY SEVIN GARDEN DUST

ACC
APR 21 1982
EPA REG. NO. 8590-257

For control of insects on vegetables, fruits, ornamentals, and lawns.

ACTIVE INGREDIENTS	
Carbaryl (1-naphthyl N-methylcarbamate)	50%
INERT INGREDIENTS	95.0%
TOTAL	100.0%

^mTrademark of Union Carbide for the active ingredient Carbaryl (1-naphthyl N-methylcarbamate)

KEEP OUT OF REACH OF CHILDREN CAUTION

See additional precautionary statements below

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid prolonged or repeated breathing of dust. Avoid skin contact. Wash hands before eating.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin, remove contaminated clothing and wash affected areas with soap and water.

If in eyes, flush eyes with plenty of water. See a physician if irritation persists.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARDS

This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION

Apply Agway Sevin Garden Dust when insects first appear preferably before damage from feeding occurs. Repeat at 7 to 10 day intervals as necessary. Apply with conventional dusting equipment. Thorough coverage to top and underside of leaves is necessary. Avoid using during flowering and pollination period. Sevin is toxic to honey bees. Do not dust too heavily or crop injury may result to tender foliage in the presence of rain or high humidity following dusting.

This product does not control mites. Where mites are a problem, use a suitable miticide.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, AND DIRECTIONS.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

VEGETABLES:

ASPARAGUS: For control of asparagus beetle and tarnished plant bug on seedlings and spears, apply 1/2 to 1/4 lbs. per 1,000 sq. ft. when insects appear and repeat as needed, but not more often than once every three days. Do not apply within 1 day of harvest.

BEANS: For control of Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetles, Leafhoppers, Western Bean Cutworm, Flea Beetles and Japanese Beetles. Apply 1/2 lb. per 1,000 sq. ft. for control of Corn Earworm, Armyworms and Velvet Bean Caterpillar. Apply 1/2 to 1/4 lb. per 1,000 sq. ft. for control of Lima Bean Pod Borer, Lygus and Stink Bugs. Apply 1 lb. per 1,000 sq. ft. May be applied up to and including the day of harvest on Beans.

CORN: For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetle, Japanese Beetle, Leafhoppers, Flea beetles and corn rootworm (adults). Apply 1/2 to 1 lb. per 1,000 sq. ft. for larvae in whorl (budworm damage) and foliage feeders; apply to entire plant. Repeat as necessary for insects attacking silks and ears, apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to day of harvest on corn.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOLMRAB: For control of Flea Beetle and Harlequin Bug. Corn Earworm, Armyworm and Cabbage Caterpillars. Apply 1/2 to 2/3 lbs. per 1,000 sq. ft. Do not apply within 3 days of harvest. Repeat at 7 to 10 day intervals as needed.

CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE (ESCAROLE), GARDEN BEETS (TOPS), HOISERADISH, KALE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFFY (TOPS), SPINACH, SWISS CHARD AND TURNIPS (TOPS): For control of Flea Beetles, Harlequin Bug and Leafhoppers apply 1/2 to 1/4 lb. per 1,000 sq. ft. when insects first appear and repeat at 5 to 7 day intervals as needed. Do not apply within 14 days of harvest when leaves or tops are used. Where only roots of Beets, Radish, Horseradish, Parsnips, Rutabagas, Salsify, and Turnips are used, do not apply within 3 days of harvest.

CUCUMBERS, SQUASH (SUMMER AND WINTER), PUMPKINS, MELONS: For control of Cucumber Beetle, Melonworm, Pickleworm, Squash Bugs, Leafhopper and Flea Beetle, apply 1/2 lb. per 1,000 sq. ft. Apply when insects appear and repeat at 10 day intervals as needed. Avoid excessive applications. Some leaf injury may occur if treatment is made when tender foliage is wet or in presence of very high humidity.

CARROTS AND LETTUCE: For control of Armyworm, tarnished plant bug, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six spotted Leafhopper, Spittlebug, Lygus and Stink Bugs. Apply 1/2 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals. Do not apply within 3 days of harvest on Head Lettuce or 14 days of harvest on Leaf Lettuce.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS: For control of Colorado Potato Beetle, Tarnished plant bug, Flea Beetle and Leafhoppers. Apply 1/2 to 1/4 lb. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus Bugs and Stink Bugs. May be applied up to and including day of harvest. Apply 1 lb. per 1,000 sq. ft.

FRUITS:

BLUEBERRIES: For control of Japanese Beetle, Cherry and Cranberry Fruitworms, apply dust thoroughly covering top and underside of leaves. Apply 1/2 lb. per 1,000 sq. ft. when insects are first noticed and repeat at 7 to 10 day intervals as necessary. May be applied up to and including the day of harvest.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, AND RASPBERRIES: For control of Japanese Beetle, Leaf Rollers, Leafhoppers and Raspberry Aphid, dust the roughly covering top and underside of the leaves. Apply 1/2 lb. per 1,000 sq. ft. when insects appear and repeat at 7 to 10 day intervals as needed. Full dust coverage is essential for good control. Do not apply within 3 days of harvest.

GRAPES: For control of Grape Berry Moth, Japanese Beetle, Grape Leaf Folder and Leafhoppers, dust thoroughly covering top and underside of leaves. Apply 1/2 to 1 lb. per 1,000 sq. ft. when insects appear and repeat at 10 day intervals as needed. May be applied up to and including the day of harvest.

STRAWBERRIES: For control of Meadow Spittle Bug, Strawberry Weevil, Leafhopper, Leaf Roller, dust thoroughly covering top and underside of leaves. Apply 1/2 lb. per 1,000 sq. ft. when insects first appear and repeat at 10 day intervals as needed. Do not apply within one day of harvest.

ORNAMENTALS:

For control of Aphids, Bagworm, Blister Beetle, Boxelder Bug, Elm Leaf Beetle, Flea Beetle, Japanese Beetle, Leafhoppers, Leaf Rollers, Periodical Cicada, Plant Bugs, Scale Insect, Crawlers, Tent Caterpillars, Exposed Thrips, dust thoroughly both top and underside of leaves. Apply at 7 to 10 day intervals throughout entire growing season, starting before serious build-up of insects occurs.

Sevin has been used at recommended rates without damage to Andromeda, Arbutus, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Camellia, Carnation, Chinese Elm, Chrysanthemum, Dahlia, Delphinium, Dogwood, Elm, Gladiolus, Hemlock, Holly, Hydrangeas, Iris, Juniper, Laurel, Lilac, Magnolia, Maidenhair Fern, Maple, Marigold, Nasturtium, Oak, Ornamental Mulberry, Petunia, Privet, Rhododendron, Rose, Rose of Sharon, Spirea, Verbena, Viburnum, Willow, Wisteria, Zinnia. Apply 1 lb. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do NOT use on Boston Ivy.

LAWNS:

For control of Chinch Bugs, Fall Armyworms, Leafhoppers, Sod Webworms (Lawn Moths), Ants, Earwigs, Fleas, Millipedes, and Mosquitoes. Apply evenly 5 lbs. dust per 1,250 sq. ft. or 1 lb. per 250 sq. ft. For control of Chinch Bugs, Fall Armyworms, Sod Webworms and Ants, apply water after application to move dust into the soil where insects frequent. Repeat in 7 to 14 days if necessary.

DOGS, CATS:

Brown dog ticks fleas. Dust liberally over all the animal and rub thoroughly into the skin. Apply also to the feet between the toes and on the legs. Apply at weekly intervals, if needed. Avoid treatment of kittens and puppies less than four weeks of age.

In or Around Dog and Cat Quarriers: Ticks, fleas. Dust liberally in and around sleeping quarters, kennels, and dog houses. See into cracks and crevices. Repeat as necessary.

DISCLAIMER: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-257 EPA Est. No. 8590-PA-1

AGWAY AGWAY INC. — CHEMICAL DIV
PO BOX 4933
SYRACUSE, NY 13221

NET WT. 1 LB. A 3/82

BEST AVAILABLE COPY