

PM-19

Reg # 8590-257

AGWAY

SEVIN GARDEN DUST

For Control of Insects on Vegetables, Fruits, Ornamentals and Lawns

ACTIVE INGREDIENT

Carbaryl (1-naphthyl N-methylcarbamate)

5.00%

INERT INGREDIENTS

95.00%

Total

100.00%

SEVIN® is the Registered Trademark of Rhone Poulenc for Carbaryl Insecticide

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional
precautionary statements

EPA Reg No 8590 257

EPA Est No 769-GA-1(A),

1386-OH-1(B)

Letter following lot number indicates

EPA Establishment location

Net Weight, 4 Pounds (1.81 kg)

ACCEPTED
MAR 20 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8590-257

Manufactured for
AGWAY Inc
P O Box 4741
Syracuse, NY 13221

96 DEC -2 7 28

REC-1 /DPD1

PRECAUTIONARY STATEMENTS
Hazards To Humans And Domestic Animals

CAUTION: May be harmful if swallowed Avoid breathing dust Avoid contact with eyes, skin or clothing Wash thoroughly after handling

GARDEN USER SAFETY REQUIREMENTS
<ul style="list-style-type: none">▶ Wear long sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product▶ Change clothing as soon as possible after use▶ Wash the outside of gloves before removing As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet▶ Do not allow children or pets to contact treated area until dust has settled

FIRST AID

If Swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger Call a physician immediately Do not induce vomiting or give anything by mouth to an unconscious person

If Inhaled, remove victim to fresh air Apply artificial respiration, if indicated

If on Skin, remove contaminated clothing and wash affected areas with soap and water

If in Eyes, flush eyes with plenty of water See a physician if irritation persists

NOTE TO PHYSICIAN AND VETERINARIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor Atropine is antidotal

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark Do not contaminate water by cleaning of equipment or disposal of wastes

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels

GENERAL INFORMATION

Apply AGWAY SEVIN Dust when insects first appear, preferably before damage from feeding occurs Repeat at 7 to 10 day intervals as necessary Apply with conventional dusting equipment Thorough coverage to top and underside of leaves is necessary Avoid using during flowering and pollination period SEVIN is toxic to honey bees Do not dust too heavily or crop injury may result to tender foliage in the presence of rain or high humidity following dusting

This product does not control mites Where mites are a problem, use a suitable miticide

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust

Apply when air is calm to avoid drift and contact with eyes and skin

Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces Allow dust to settle in treated areas before reentering

For plants above chest height, consider using a liquid carbaryl product

Do not repeat application more than once a week

Vegetables

ASPARAGUS For control of asparagus beetle and tarnished plant bug on seedlings and spears apply 1/2 to 3/4 lbs per 1 000 sq ft when insects appear and repeat as needed but not more often than once every three days Do not apply within 1 day of harvest

BEANS For control of Mexican bean beetle bean leaf beetle, cucumber beetles leafhoppers western bean cutworm flea beetles and Japanese beetles apply 1/2 lb per 1 000 sq ft For control of corn earworm armyworms and velvet bean caterpillar apply 1/2 to 3/4 lb per 1 000 sq ft For control of lima bean pod borer lygus and stink bugs apply 1 lb per 1 000 sq ft May be applied up to and including the day of harvest on beans

CORN For control of corn earworm European corn borer fall armyworm sap beetle Japanese beetle, leafhoppers flea beetles and corn rootworm (adults) apply 1/2 to 1 lb per 1 000 sq ft For larvae in whorl (budworm damage) and foliage feeders apply to entire plant Repeat as necessary For insects attacking silks and ears apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry Three or more applications may be required depending on severity of infestation Timing and good coverage are absolutely essential for effective control May be applied up to day of harvest on corn

BEST AVAILABLE COPY

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI For control of flea beetle and harlequin bug, corn earworm, armyworm and cabbage caterpillars, apply 1/2 to 2/3 lbs per 1 000 sq ft. Do not apply within 3 days of harvest. Repeat at 7 to 10 day intervals as needed.

CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE (ESCAROLE), GARDEN BEETS (TOPS), HORSE RADISH, KALE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY (TOPS), SPINACH, SWISS CHARD AND TURNIPS (TOPS) For control of flea beetles, harlequin bug and leafhoppers, apply 1/2 to 3/4 lb per 1 000 sq ft when insects first appear and repeat at 5 to 7 day intervals as needed. Do not apply within 14 days of harvest when leaves or tops are used. Where only roots of beets, radish, horseradish, parsnips, rutabagas, salsify and turnips are used, do not apply within 3 days of harvest.

CUCUMBERS, SQUASH (SUMMER AND WINTER), PUMPKINS, MELONS For control of cucumber beetle, melonworm, pickleworm, squash bugs, leafhopper and flea beetle, apply 1/2 lb per 1 000 sq ft. Apply when insects appear and repeat at 10 day intervals as needed. Avoid excessive applications. Some leaf injury may occur if treatment is made when tender foliage is wet or in presence of very high humidity.

CARROTS AND LETTUCE For control of armyworm, tarnished plant bug, cabbage caterpillar, small cabbage looper, corn earworm, flea beetle, harlequin bug, leafhopper, spotted leafhopper, spittlebug, lygus and stink bugs, apply 3/4 lb per 1 000 sq ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals. Do not apply within 3 days of harvest on head lettuce or 14 days of harvest on leaf lettuce.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS For control of Colorado potato beetle, tarnished plant bug, flea beetle and leafhoppers, apply 1/2 to 3/4 lb per 1 000 sq ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, lygus bugs and stink bugs, may be applied up to and including day of harvest. Apply 1 lb per 1 000 sq ft.

Fruits

BLUEBERRIES For control of Japanese beetle, cherry and cranberry fruitworms, apply dust thoroughly covering top and underside of leaves. Apply 3/4 lb per 1 000 sq ft when insects are first noticed and repeat at 7 to 10 day intervals as necessary. May be applied up to and including the day of harvest.

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, AND RASPBERRIES For control of Japanese beetle, leafrollers, leafhoppers and raspberry aphid, dust thoroughly covering top and underside of the leaves. Apply 3/4 lb per 1,000 sq ft when insects appear and repeat at 7 to 10 day intervals as needed. Full dust coverage is essential for good control. Do not apply within 7 days of harvest.

GRAPES For control of grape berry moth, Japanese beetle, grape leaf folder and leafhoppers, dust thoroughly covering top and underside of leaves. Apply 3/4 to 1 lb per 1 000 sq ft when insects appear and repeat at 10 day intervals as needed. May be applied up to and including the day of harvest.

STRAWBERRIES For control of meadow spittlebug, strawberry weevil and strawberry leafroller, dust thoroughly covering top and underside of leaves. Apply 3/4 lb per 1 000 sq ft when insects first appear and repeat at 10 day intervals as needed. Do not apply within one day of harvest.

BEST AVAILABLE COPY

Ornamentals

For control of aphids, bagworm, blister beetle, boxelder bug, elm leaf beetle, flea beetle, Japanese beetle, leafhoppers, leafrollers, periodical cicada, plant bugs, scale, insect crawlers, tent caterpillars, exposed thrips. Dust thoroughly both top and underside of leaves. Apply at 7 to 10 day intervals throughout entire growing season, starting before serious build up of insects occurs.

SEVIN has been used at recommended rates without damage to andromeda, arbutus, ash, aster, azalea, beech, begonia, birch, boxwood, camellia, carnation, Chinese elm, chrysanthemum, delphinium, dogwood, elm, gladiolus, hellebore, holly, hydrangeas, jasminum, laurel, lilac, magnolia, mandarin, maple, marigold, nasturtium, oak, ornamental mulberry, petunia, privet, rhododendron, rose, rose of Sharon, spirea, verbena, viburnum, willow, wisteria, zinnia. Apply 1 lb per 1000 sq ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals as necessary. Do NOT use on Boston ivy.

Lawns

For control of chinch bugs, fall armyworms, leafhoppers, sod webworms (lawn moths), ants, earwigs, fleas, millipedes and mosquitoes. Apply evenly 5 lbs dust per 1250 sq ft or 1 lb per 250 sq ft. For control of chinch bugs, fall armyworms, sod webworms and ants, apply water after application to move dust into the soil where insects frequent. Repeat at 7 to 14 days if necessary.

**READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS OR CATS**

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

PET USER SAFETY REQUIREMENTS

- ▶ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product
- ▶ Change clothing as soon as possible after use
- ▶ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

Use AGWAY SEVIN Dust only outside or in a well ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Reapply every week if needed. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Consult a veterinarian before using this product on medicated, debilitated, aged, or nursing animals. Dust bedding and doghouse using a shaker can or other dust applicator.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

DISCLAIMER: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

NOTE: This pesticide is to sold ONLY in this original unbroken package.

FOR PRODUCT INFORMATION CALL: (315)449-6244