



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

Beverly Randall  
Agway Inc.  
c/o Universal Cooperatives, Inc.  
P.O. Box 460  
Minneapolis, MN 55440

MAR 2 1998

Subject: Sevin Garden Dust  
EPA Registration No. 8590-257  
Deletion of Trees from Label  
Amendment dated March 4, 1997  
Federal Register (Vol. 62, No. 166) August 27, 1997

Dear Ms. Randall:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you:

1. Make the following changes to your label under the heading "VEGETABLES":
  - a) In the Directions for Use for CORN, change the application interval from "2 to 3 days" to "apply at 7 day intervals".
  - b) In the Directions for Use on CHINESE CABBAGE, etc.... change the "repeat at 5 to 7 day intervals.." to "repeat at 7 day intervals as needed".
  - c) In the Directions for Use for CARROTS AND LETTUCE change the "repeat as needed at 5 to 7 day intervals" to "repeat as needed at 7 day intervals".
2. Submit two copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions. A copy of the labeling stamped "Accepted with Comments" is enclosed for your records.

Sincerely,

*George Tompkins*

George Tompkins, Ph.D., Entomologist  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

297

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

MAR 2 1998

AGWAY

SEVIN GARDEN DUST

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
8590-257

For Control of Insects on Vegetables, Fruits, and Lawns

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl-N-methylcarbamate) . . . . .	5.00%
INERT INGREDIENTS: . . . . .	95.00%
Total . . . . .	100.00%

SEVIN® is the Registered Trademark of Rhone Poulenc for Carbaryl Insecticide.

KEEP OUT OF REACH OF CHILDREN  
CAUTION

See back panel for additional  
precautionary statements.

REC'D  
97 MAR 11 12:11  
/DPD1

EPA Reg. No. 8590-257  
EPA Est. No. 769-GA-1(A),  
1386-OH-1(B)  
Letter following lot number indicates  
EPA Establishment location.

Net Weight: 4 Pounds (1.81 kg)

Manufactured for:  
AGWAY Inc.  
P.O. Box 4741  
Syracuse, NY 13221

**PRECAUTIONARY STATEMENTS**  
**Hazards To Humans And Domestic Animals**

**CAUTION:** May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

GARDEN USER SAFETY REQUIREMENTS
<ul style="list-style-type: none"> <li>▶ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.</li> <li>▶ Change clothing as soon as possible after use.</li> <li>▶ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.</li> <li>▶ Do not allow children or pets to contact treated area until dust has settled.</li> </ul>

**FIRST AID**

**If Swallowed,** drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

**If Inhaled,** remove victim to fresh air. Apply artificial respiration, if indicated.

**If on Skin,** remove contaminated clothing and wash affected areas with soap and water.

**If in Eyes,** flush eyes with plenty of water. See a physician if irritation persists.

**NOTE TO PHYSICIAN AND VETERINARIAN:** Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

**ENVIRONMENTAL HAZARDS**

This product is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.**

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

**GENERAL INFORMATION**

Apply AGWAY SEVIN Garden Dust when insects first appear, preferably before damage from feeding occurs. Repeat at 7 to 10 day intervals as necessary. Apply with conventional dusting equipment. Thorough coverage to top and underside of leaves is necessary. Avoid using during flowering and pollination period. SEVIN is toxic to honey bees. Do not dust too heavily or crop injury may result to tender foliage in the presence of rain or high humidity following dusting.

This product does not control mites. Where mites are a problem, use a suitable miticide.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.

Apply when air is calm to avoid drift and contact with eyes and skin.

Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

For plants above chest height, consider using a liquid carbaryl product.

Do not repeat application more than once a week.

**Vegetables**

**ASPARAGUS:** For control of asparagus beetle and tarnished plant bug on seedlings and spears, apply 1/2 to 3/4 lbs. per 1,000 sq.ft. when insects appear and repeat as needed, but not more often than once every three days. Do not apply within 1 day of harvest.

**BEANS:** For control of Mexican bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, western bean cutworm, flea beetles and Japanese beetles: apply 1/2 lb. per 1,000 sq.ft. For control of corn earworm, armyworms and velvet bean caterpillar: apply 1/2 to 3/4 lb. per 1,000 sq.ft. For control of lima bean pod borer, lygus and stink bugs, apply 1 lb. per 1,000 sq.ft. May be applied up to and including the day of harvest on beans.

**CORN:** For control of corn earworm, European corn borer, fall armyworm, sap beetle, Japanese beetle, leafhoppers, flea beetles and corn rootworm (adults): apply 1/2 to 1 lb. per 1,000 sq.ft. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to day of harvest on corn.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRABI:** For control of flea beetle and harlequin bug, corn earworm, armyworm and cabbage caterpillars, apply 1/2 to 2/3 lbs. per 1,000 sq.ft. Do not apply within 3 days of harvest. Repeat at 7 to 10 day intervals as needed.

**CHINESE CABBAGE, COLLARDS, DANDELION, ENDIVE (ESCAROLE), GARDEN BEETS (TOPS), HORSERADISH, KALE, MUSTARD GREENS, PARSLEY, PARSNIPS, RADISHES, RUTABAGAS, SALSIFY (TOPS), SPINACH, SWISS CHARD AND TURNIPS (TOPS):** For control of flea beetles, harlequin bug and leafhoppers, apply 1/2 to 3/4 lb. per 1,000 sq.ft. when insects first appear and repeat at 5 to 7 day intervals as needed. Do not apply within 14 days of harvest when leaves or tops are used. Where only roots of beets, radish, horseradish, parsnips, rutabagas, salsify, and turnips are used, do not apply within 3 days of harvest.

**CUCUMBERS, SQUASH (SUMMER AND WINTER), PUMPKINS, MELONS:** For control of cucumber beetle, melonworm, pickleworm, squash bugs, leafhopper and flea beetle, apply 1/2 lb. per 1,000 sq.ft. Apply when insects appear and repeat at 10 day intervals as needed. Avoid excessive applications. Some leaf injury may occur if treatment is made when tender foliage is wet or in presence of very high humidity.

**CARROTS AND LETTUCE:** For control of armyworm, tarnished plant bug, cabbage caterpillar, small cabbage looper, corn earworm, flea beetle, harlequin bug, leafhopper, sixspotted leafhopper, spittlebug, lygus and stink bugs, apply 3/4 lb. per 1,000 sq.ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals. Do not apply within 3 days of harvest on head lettuce or 14 days of harvest on leaf lettuce.

**POTATOES, TOMATOES, EGGPLANTS AND PEPPERS:** For control of Colorado potato beetle, tarnished plant bug, flea beetle and leafhoppers: apply 1/2 to 3/4 lb. per 1,000 sq.ft. For control of tomato fruitworm, fall armyworm, tomato hornworm, European corn borer, lygus bugs and stink bugs: may be applied up to and including day of harvest. Apply 1 lb. per 1,000 sq.ft.

**Fruits**

**BLUEBERRIES:** For control of Japanese beetle, cherry and cranberry fruitworms, apply dust thoroughly covering top and underside of leaves. Apply 3/4 lb. per 1,000 sq.ft. when insects are first noticed and repeat at 7 to 10 day intervals as necessary. May be applied up to and including the day of harvest.

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, AND RASPBERRIES:** For control of Japanese beetle, leafrollers, leafhoppers and raspberry aphid, dust thoroughly covering top and underside of the leaves. Apply 3/4 lb. per 1,000 sq.ft. when insects appear and repeat at 7 to 10 day intervals as needed. Full dust coverage is essential for good control. Do not apply within 7 days of harvest.

**GRAPES:** For control of grape berry moth, Japanese beetle, grape leaf folder and leafhoppers, dust thoroughly covering top and underside of leaves. Apply ¾ to 1 lb. per 1,000 sq.ft. when insects appear and repeat at 10 day intervals as needed. May be applied up to and including the day of harvest.

**STRAWBERRIES:** For control of meadow spittlebug, strawberry weevil and strawberry leafroller, dust thoroughly covering top and underside of leaves. Apply ¾ lb. per 1,000 sq.ft. when insects first appear and repeat at 10 day intervals as needed. Do not apply within one day of harvest.

**LAWNS**

For control of chinch bugs, fall armyworms, leafhoppers, sod webworms (lawn moths), ants, earwigs, fleas, millipedes, and mosquitoes. Apply evenly 5 lbs. dust per 1,250 sq.ft. or 1 lb. per 250 sq.ft. For control of chinch bugs, fall armyworms, sod webworms and ants, apply water after application to move dust into the soil where insects frequent. Repeat in 7 to 14 days if necessary.

**READ ENTIRE LABEL BEFORE EACH USE  
USE ONLY ON DOGS OR CATS**

Do not use this product on animals simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals.

<b>PET USER SAFETY REQUIREMENTS</b>
<ul style="list-style-type: none"> <li>▶ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.</li> <li>▶ Change clothing as soon as possible after use.</li> <li>▶ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.</li> </ul>

Use AGWAY SEVIN Garden Dust only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Reapply every week if needed. This product should not be applied by children. **Do not use on pregnant dogs.** Do not use on kittens or puppies under 12 weeks of age. Consult a veterinarian before using this product on medicated, debilitated, aged, or nursing animals. Dust bedding and doghouse using a shaker can or other dust applicator.

Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

**DISCLAIMER:** Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

**NOTE:** This pesticide is to sold ONLY in this original unbroken package.

**FOR PRODUCT INFORMATION CALL: (315)449-6244**