

AGWAY
50% SEVIN

AGWAY

NOTE TO PHYSICIAN: Carbaryl* is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

*Trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate. U. S. Pat. Nos. 2,903,478 and 3,009,855.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

MANUFACTURED BY AGWAY INC.
CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201
USDA Reg. No. 8590-266

50% sevin* WETTABLE POWDER

FOR CONTROL OF INSECTS
ON ORNAMENTALS, FRUITS
AND VEGETABLES

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl
N-methylcarbamate) 50%

INERT INGREDIENTS:

(Not more than) 50%

Total 100%

caution:

**KEEP OUT OF REACH OF
CHILDREN**

See additional cautions and warnings on side panel.

NET WEIGHT 14 Oz.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of dust or spray. Do not take internally. Skin contact with dust or spray material may be harmful. Avoid contact with skin. Wash thoroughly after using. Wash hands and face before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothes daily. Avoid contamination of food, feeding troughs and watering receptacles. Keep children and pets off treated areas until insecticide is washed into soil and grass is thoroughly dry.

This product is toxic to fish. Do not contaminate any body of water. Do not reuse empty container. Destroy it by burying with wastes or burning. Stay away from smoke.

SP1068

directions

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

FOR CONTROL OF INSECTS ON: Ornamentals, Lawn Insects, Apples and Pears, Blackberries, Boysenberries, Dewberries, Loganberries and Raspberries, Blueberries, Cranberries, Grapes, Peaches, Apricots, Nectarines, Plums, Prunes, Cherries, Strawberries, Asparagus, Beans, Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi, Carrots, Chinese Cabbage, Collards, Dandelions, Endive (Escarole), Garden Beets (tops), Horseradish, Kale, Mustard Greens, Parsley, Parsnips, Radishes, Rutabagas, Salsify (tops), Spinach, Swiss Chard and Turnips (tops), Corn, Cucumbers, Squash (Summer and Winter), Pumpkins, Melons, Lettuce, Potato, Tomato, Eggplant and Pepper.

RATE OF APPLICATION: Is based on 1 tablespoonful of Agway 50% Sevin per gallon of water is equal to 1 pound per acre or 1 pound per 100 gallons of spray.

TIMING OF APPLICATION: Development of insects vary widely depending on local conditions. Good results will be obtained if sprays are applied at regular intervals when insect first appears and before serious damage occurs.

NOTE: Agway 50% Sevin does not control mites but is compatible with all common miticides. This material is unstable under highly alkaline conditions. Do not use with lime, Bordeaux, lime sulfur, casein-lime spreaders or other alkaline materials. SEVIN is toxic to honeybees. Avoid using during flowering and pollination period. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following spraying.