

Leafhopper, spittlebug	1 to 2 qts.
Alfalfa caterpillar, flea beetles	1½ to 2 qts.
Alfalfa weevil (larvae), alfalfa webworm, fall armyworm, clover leaf weevil, Mexican bean beetle, pea weevil, soybean caterpillar, velvet bean caterpillar	2 to 3 qts.
Cowpea curculio, Japanese beetle	3 qts.
Armyworm	3 to 4 qts.

DO NOT APPLY WITHIN 7 DAYS OF HARVEST OR GRAZING.

POTATOES — (Colorado potato beetle, flea beetles, leafhoppers): Use 2 qts. of Methoxychlor 2E per acre in 100 gallons of water.

GRAIN STORAGE BINS — To clean up infestations in empty storage bins, use 1 gal. Methoxychlor 2-E in 10 gals. of water applied at the rate of 2 gals. of spray per 1,000 sq. ft. before products are stored. This aids in control of cadelle, flat grain beetle, sawtoothed grain beetle, lesser grain borer, red toothed grain beetle, red flour beetle, long-headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle.

NOTE — Do not add grain to bin for at least 24 hours or until walls have dried out thoroughly.

CONTACT AND SPACE SPRAY FOR FLIES — To reduce population of houseflies and stable flies in farm buildings, in barnyards, in alleys, on dumps and other outdoor locations, dilute 1 pt. of Methoxychlor 2-E with 2½ gals. of water and apply in space by mist blower or directly to flies by hydraulic sprayer, hand sprayer or cylindrical pressure sprayer. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying.

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. Avoid breathing of spray mist. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply when weather conditions favor drift of spray from areas treated.

This product is toxic to fish. Keep out of any body of water.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes, or by burning (keep out of smoke).

IMPORTANT—Users should observe the usual precautions concerning application of oil-containing products to crops during high temperatures, since many crops are then more susceptible to oil injury. IF EXPOSED TO PROLONGED COLD, PLACE IN WARM STORAGE (50° to 80° F.) FOR SEVERAL HOURS AND AGITATE BEFORE USING.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. IF EXPOSED TO PROLONGED COLD, PLACE IN WARM STORAGE (50° to 80° F.) FOR SEVERAL HOURS AND AGITATE BEFORE USING.

DISCLAIMER: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

EPA Est. No. 8590—PA.—1

EPA Reg. No. 8500-267

267

**AGWAY INC.
FERTILIZER — CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201**

NET CONTENTS FIVE GALLONS

