

CLFANS

RESTORES

MAKES FLOORS MORE CONDUCTIVE

FOR ALL SURFACES INCLUDING TERRAZZO, ASPHALT TILE, VINYL, RUBBER, LINOLEUM, MARBLE OR CORK.

THE ECONOMICAL WAY TO CLEAN, DISINFECT, DEODORIZE AND DELIME the following types of Surfaces: Floor — including Terrazzo, Ceramic Tile, Vitreous Tile, Asphalt Tile, Rubber, Vinyl Linoleum, Magnesite, Concrete, etc. Painted, Enameled, Varnished, and Lacquered Surfaces, including Walls, Woodwork, Cabinets, Refrigerators, Display Cases, Fixtures and Utensils.

When Quatra-D is used in Dairies, Food Processing Plants, Restaurants and Taverns on utensils, glasses, dishes and food handling equipment, rinse thoroughly with potable water, either immediately after treatment of the surfaces with Quatra-D, or just prior to reuse.

Cleans more rapidly and surface stays cleaner longer. May be used even in hard water.

Removes Soap Scum and Alkaline Salts.

Eliminates odors quickly because of fixation and destruction of bacteria at the source. LEAVES NO AFTER ODOR.

GENERAL DIRECTIONS

FOR FLOORS, Use 4 oz. per gallon of Lukewarm Water and Mop or Scrub in usual manner. In this dilution it will clean a waxed floor without removing the wax. FOR WALLS, Furniture, Fixtures, and other Varnished, Waxed, or Enameled Surfaces. Use 3 ounces per gallon of warm water.

Used extensively in Hospitals, Institutions, and Industry

IT DISINFECTS

as it DEODORIZES

as it SANITIZES

DIRECTIONS

AS A DISINFECTANT: After Surface has been thoroughly cleaned, use 2 oz. to each gallon of water for non-porous surfaces and at least 4 oz. to each gallon of water for porous surfaces.

FOR DEODORIZING AND SANITIZING: Use 1 oz. per gallon of water (23/4 ppm).

When Quatra-D is used as a sanitizer for equipment and utensils in official meat and poultry processing establishments, the maximum allowable concentration is 200 ppm, or approximately 1 3/4 ounces of Quatra-D per 2 gallons of water. Rinse thoroughly with potable water, either immediately after treatment of the surfaces with Quatra-D, or just prior to reuse.

Germicidal activity will be impaired with Soap or other Detergents.

Guarantee

We hereby guarantee this Disinfectant to have a Phenol Coefficient of 25 when tested against antibiotic resistant strain of Staphylococcus Aureus Phage Type 80. Phenol Coefficient 15 against S. typhosa.

NON-FLAMMABLE

CAUTION: Avoid contamination of food. Avoid getting the concentrate in eyes or on the skin. In case of contact, flush with plenty of water. If irritation persists, get medical attention.

33-11/67 2M LL

AGWAY PROMOTIONAL SERVICES

COPY

FOR: AGWAY SANITIZER

DATE: 1/10/73

((FRONT PANEL))

P.C.A.—50 CHARACTERS
E.U.T.—80 CHARACTERS

((AGWAY T.M.))

NET CONTENTS
ONE GALLON

SANITIZER

((FIVE GALLON
SIZE ALSO))

For Hatchery, Poultry and Livestock
Premise Disinfection

ACTIVE INGREDIENTS:

n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈) dimethyl benzyl ammonium chlorides.....	10.0%
n-Alkyl (50% C ₁₂ , 30% C ₁₄ , 17% C ₁₆ , 3% C ₁₈) dimethyl ethylbenzyl ammonium chlorides.....	10.0%

INERT INGREDIENTS:.....80.0%

100.0%

DANGER: ((18 pt. type)) KEEP OUT OF REACH OF
CHILDREN. ((12 pt. type)) See additional
instructions and precautions on side panel.
REAR

MANUFACTURED BY AGWAY INC.
Fertilizer-Chemical Division
Box 1333
Syracuse, N. Y. 13201

EPA Reg. No.
8590-270

OK BY

ACCEPTED

FILED

UNDER

FOR

ED

AGWAY PROMOTIONAL SERVICES

COPY

FOR: AGWAY SANITIZER

DATE: 1/10/75
20/16/75

((REAR PANEL))

PICA—50 CHARACTERS
ELITE—60 CHARACTERS

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONS AND DIRECTIONS.

For general farm use, in hatcheries, for dairy use or for use in pig parlors dilute 1 1/2 ounces (3 tablespoonfuls) Agway Sanitizer to .4 gallons of water. This supplies 500 ppm of active quaternary sanitizer. 526

O.K. BY

ACCEPTED
FEB 10 1975
UNITED STATES DEPARTMENT OF AGRICULTURE
FOR ECONOMIC ENTOMOLOGICAL SERVICE
ED UNDER NO. 110-1-100

20F4
20/16/75

AGWAY PROMOTIONAL SERVICES

COPY

FOR: AGWAY SANITIZER

DATE: 1/10/73
SDF/ST/RS

((REAR PANEL)) continued

PC-30 CHARACTERS
ELITE-60 CHARACTERS

FOLLOW THESE GENERAL USE PROCEDURES:

- a. Remove all poultry, ^{LIVESTOCK,} and feeds from premises, cars, trucks, coops, crates.
- b. Remove all litter and droppings from floors, walls, surfaces of facilities and fixtures traversed or occupied by poultry ^{OR LIVESTOCK.}
- c. Empty all troughs, racks, and other feedings and watering appliances.
- d. Saturate all surfaces with disinfectant solution.
- e. Immerse all forks, shovels, and scrapers.
- f. Ventilate buildings, cars, boats, coops, and other closed spaces. Do not house ^{LIVESTOCK} poultry, or employ equipment until treatment has been absorbed, set, or dried.
- g. All treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be scrubbed thoroughly with detergents and rinsed with potable water prior to reuse.

ADVANTAGES

Provides positive deodorization of treated surfaces.

Kills Mycoplasma gallisepticum (CRD) to reduce the hazard of cross contamination to environmental surfaces during outbreaks.

ACCEPTED

FOR THE USE OF THE UNITED STATES GOVERNMENT

ED UNDER NO. 1

30F4