

ACCEPTED

APR 11 1967

AGWAY  
T.M.

Net Weight 4 Pounds

AGWAY CORN HERBICIDE

For Selective Weed Control in Corn

ACTIVE INGREDIENTS:

|  |        |
|--|--------|
| Linuron /3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea/ .....  | 30.8%  |
| Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine) . | 30.9%  |
| INERT INGREDIENTS: .....                                       | 38.4%  |
| TOTAL .....  | 100.0% |

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

CAUTION: (18 pt. type) KEEP OUT OF REACH OF CHILDREN. (12 pt. type) Harmful if swallowed. May irritate eyes, nose, throat, and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Avoid contamination of feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of any body of water.

Do not apply when conditions favor drift or runoff from the crop area treated. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning (Keep out of smoke.).

Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas.

Apply this product only as specified on this label.

STATEMENT OF LIABILITY: The use of this material being beyond our knowledge and control we make no representation or warranty, express or implied, concerning this material except that it conforms to the chemical description on the label. Use of this material in a manner or at a time other than in accordance with directions may produce plant injury. We shall not be responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage, or use of this material. The buyer accepts and uses this material on these conditions.

DIRECTIONS FOR USE

Make a single application as a band (over the row) or broadcast spray after planting but before crop emerges. For accuracy and convenience, planting and spraying should be combined in one operation. Apply with a properly calibrated fixed-boom power sprayer, using sufficient water (equivalent to 25 to 40 gals. per acre on a broadcast basis) to provide thorough and uniform coverage of the ground. Avoid overlapping, and shut off spray boom while starting, turning, slowing or stopping, or injury to the crop may result. Continuous agitation is required to keep the material in suspension.

Best results are obtained when Agway Corn Herbicide is moved by moisture (rainfall or overhead irrigation) into the upper layer of soil within 2 weeks

after application. Lack of sufficient moisture (1 to 2 inches is generally sufficient) following treatment will generally result in unsatisfactory weed control.

Soil should be well prepared and as free as possible from trash and clods. Seed should be planted to a depth of at least 1 3/4" or crop injury may result. If soil becomes crusted before corn emerges, a shallow rotary hosing may be made. If rainfall is insufficient following treatment to activate the chemical, rotary hosing or standard cultivation should be used before weeds become well established; otherwise, avoid cultivating or disturbing the sprayed area, such as covering treated band with soil while cultivating middles.

Agway Corn Herbicide is a wettable powder to be mixed in water and applied as a spray for the selective control of annual weeds and grasses in corn. It is non-corrosive, non-volatile and non-flammable.

Treatment provides control of germinating annual weed seedlings such as foxtail (including giant foxtail), barnyardgrass, fall panicum, crabgrass, buttonweed, lambsquarters, pigweed, purslane, mustard, ragweed, and smartweed. Treatment will not control established nutgrass, Johnsongrass, Bermudagrass, or other perennial weeds.


Apply the spray mixture at the following rates:

| Soil Type*  | Lbs. Agway Corn Herbicide in 5 to 40 gals. water per acre (of ground actually sprayed) | Example -- Band Treatment Corn Herbicide/crop acre 14" Band on 42" Rows |
|---|--|---|
| <b>Light colored -- low organic matter (2% or less)</b> |  |   |
| Sandy loam  | 1.6  | .5  |
| Silt loam   | 2.0  | .7  |
| Clay loam   | 2.5  | .8  |
| <b>Dark colored -- moderate organic matter</b>          |  |   |
| Silt loam   | 4.0  | 1.3   |
| Clay loam   | 5.0  | 1.7   |

\*Do not use on sand or loamy sand soil as crop injury may result.

Fields treated with Agway Corn Herbicide may be replanted to the same crop if initial seeding fails to germinate or produce a stand. Thoroughly rework soil before replanting. Do not retreat field with a second application during the same crop year as injury to the crop may result. Do not replant treated areas with crops other than corn within 5 months after application or crop injury may result. Do not follow treated corn with sugar beets, tobacco or vegetables in rotation. Prior to replanting to any crop following treated corn, thorough seedbed preparation including fall or spring plowing is recommended.

EQUIPMENT: Use a tractor-mounted power sprayer properly calibrated to a constant speed and rate of delivery (see CALIBRATION section). Sprayer should be designed to treat the same number of rows as the planter used. Make certain



spray equipment is clean (free of scale, rust, dirt, oil, and pesticide deposits). Screens should be 50-mesh or coarser. Booms and hoses should be maximum of 1/2" diameter. MATERIAL MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION. Agitate by mechanical or hydraulic means. With hydraulic system, use a separate agitation line in addition to by-pass line; both lines should terminate at bottom of tank to minimize foaming. Spray booms must be shut off while starting, turning, slowing or stopping, or injury to the crop may result.

**CALIBRATION:** The following is designed for application of indicated number of gals. spray per acre of ground actually sprayed; it is based on width of band treated and is not affected by row spacing.

**Step 1:** To calibrate, use tractor gear and throttle setting which will be used for actual application (full throttle is preferred to insure maximum pump capacity). With only water in the tank, determine number of seconds required for tractor to travel in the field the measured distance shown below -- opposite desired band width and under proper spray volume.

| Band Width | Gals. Spray Per Acre of Ground Actually Sprayed |         |         |         |
|------------|---|---------|---------|---------|
|            | 25  | 30      | 35      | 40      |
| 10 in.     | 523 ft.   | 436 ft. | 363 ft. | 326 ft. |
| 12 in.     | 436 ft.   | 363 ft. | 311 ft. | 272 ft. |
| 14 in.     | 374 ft.   | 311 ft. | 266 ft. | 232 ft. |
| 16 in.     | 327 ft.   | 272 ft. | 233 ft. | 204 ft. |
| 18 in.     | 291 ft.   | 242 ft. | 207 ft. | 181 ft. |
| 20 in.     | 262 ft.   | 218 ft. | 187 ft. | 163 ft. |

Adjust height of boom to obtain desired band width. With only water in the tank and using the same throttle setting as in Step 1 but with tractor standing still, adjust pressure so that EACH NOZZLE delivers ONE QUART of spray in the exact number of seconds as determined in Step 1 for the selected band width.

**Step 2:** Equipment is now ready to apply the required spray volume, provided nozzle height, gear and throttle setting are not changed.

**NOTE:** This pesticide is to be sold ONLY in this original, unbroken package.

MANUFACTURED BY: Agway, Inc., Syracuse, N.Y.

USDA Reg. No. 8590-

