

CHLORDANE 25G

For the Control of Soil Insects in Corn (Field and Sweet)

ACCEPTED

Oct 23-57

THE FEDERAL
AGRICULTURAL
PESTICIDE AND
PLANT QUARANTINE
SERVICE
ECONOMIC INSECTICIDES
DIVISION
UNDER NO. 8590

NET WEIGHT FIFTY POUNDS

ACTIVE INGREDIENTS:

Technical Chlordane*	25.0%
INERT INGREDIENTS	75.0%
	100.0%

*Equivalent to 15% Octachloro-4, 7-methano-tetrahydroindane and 10% related compounds.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust and spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

ANTIDOTE: External—In case of spillage on skin, wash promptly with soap and water. Internal—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

Container Disposal: This container must not be reused. Bury unused contents 18 inches beneath the soil surface away from water and sites of planned excavation. Burn a few containers at a time to avoid large fires. Stay away from smoke.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

All lb. per acre dosage rates mentioned in the following directions are to be used for applying Chlordane 25 G as it is formulated in this package.

FIELD CROPS: For use on Corn (Field and Sweet), Peanuts and Tobacco.

Broadcast application: Apply 10 lbs. of Chlordane 25 G per acre for control of Northern, Southern, and Western corn rootworms, corn field ants, and corn root aphids. Apply 10-20 lbs. per acre for control of the larvae of seed corn beetles, Asiatic garden beetles, green June bugs, and white fringed beetles. Apply 40 lbs. per acre to control Japanese beetle larvae. For seed corn maggot, cutworm, white grub, and wireworm control, apply 10-16 lbs. per acre. Apply Chlordane 25 G after the field is plowed. Disc in promptly.

Use the higher rate on heavier soils and the lower rate on light, sandy soils. Use the higher rates for heavy infestation of cutworms.

Row Application: For the control of Northern, Southern, and Western corn rootworm, corn field ants, and corn root aphids, apply 4 lbs. of Chlordane 25 G per acre. The seed corn maggot, cutworms, white grubs, and wireworms, as well as larvae of the seed corn beetle, may be controlled by an application of 8 lbs. per acre. Apply Chlordane 25 G at planting time in the furrow, behind the planter shoe and ahead of the press wheel.

VEGETABLE CROPS: On Beans, Beets, Blackeyed Peas, Broccoli, Brussel Sprouts, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cowpeas, Cucumbers, Eggplant, Kale, Kohlrabi, Eggplant, Melons, Okra, Onions, Peas, Peppers, Pimientos, Potatoes, Radishes, Rutabagas, Squash, Sweet Potatoes, Tomatoes, Turnips, and Watermelons.

Broadcast application: For the control of cutworms, wireworms, and white grubs, apply 10-16 lbs. per acre. Larvae of the Asiatic garden beetle, white fringed beetle, and flea beetles (Potatoes and Sweet Potatoes) may be controlled with 10-20 pounds per acre. For control of Japanese beetle larvae, apply 40 lbs. per acre.

Row Application: Apply 8 lbs. of 25 G Chlordane per acre for control of cutworms, wireworms, and white grubs.

FRUIT CROPS: Apples, Apricots, Cherries, Nectarines, Peaches, Pears, Plums, Prunes, Quinces, Blackberries, Blueberries, Grapes, Huckleberries, Loganberries, Raspberries, and Strawberries.

Broadcast application: Apply 6-8 lbs. per acre for control of ants. Apply 10-16 lbs. per acre for the control of white grubs, cutworms, and wireworms. Apply 40 lbs. per acre for control of Japanese beetle larvae, Larvae of the Asiatic garden beetle, green June bug, and white fringed beetle may be controlled with 10-20 lbs. per acre. Blackberries, Blueberries, Raspberries, and Strawberries, use 20-40 lbs. per acre to control root weevil larvae and rootworms. On berry crops make application at base of plants and incorporate into soil. Do not apply when fruit is present.

ORNAMENTALS: On ornamental turf, lawns, nursery and greenhouse soils.

Broadcast application: To control ants, ticks and chiggers, apply 6 lbs. per acre. Crickets, European earwigs, grasshoppers, adult mosquitoes, rose chafers and spiders may be controlled by applying 10-20 lbs. per acre. Apply 20-40 lbs. per acre for the control of white grubs, wireworms, and larvae of the Asiatic garden beetle, European chafer, green June bug and white fringed beetle. Apply 40 lbs. per acre for control of Japanese beetle larvae.

On turf, lawns, nursery and greenhouse soils, apply granules uniformly, using a lawn fertilizer spreader or suitable granular equipment. On bare soil incorporate the application into the soil immediately. Birds feeding on treated areas may be harmed.

Do not allow children, pets or domestic animals on treated lawn until the insecticide is washed off the grass into the soil by sprinkling or rainfall and grass is completely dry.

Consult your State Agricultural Extension Service for detailed recommendations.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

USDA Reg. No. 8590-273

AC-273G

Manufactured by AGWAY INC.
Chemical Division
Box 1333, Syracuse, N.Y. 13201