

U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF PLANT HEALTH AND
QUARANTINE
CAPTAIN FOR CONTROL
OF INSECTS AND DISEASES
ON VEGETABLES

CAUTION:


KEEP OUT OF REACH
OF CHILDREN

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

MANUFACTURED BY AGWAY INC.
CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201

USDA Reg. No. 8590-283 11-68 Tr



CAUTION: Harmful if inhaled, swallowed or absorbed through skin. Avoid skin contact with dust. Avoid breathing of dust. Do not take internally. Wash thoroughly after using and before eating. Avoid contamination of feed and food-stuffs. Do not over-dose. Avoid use on wet foliage or in presence of high humidity or high temperature of several days duration.

This product is toxic to fish. Do not contaminate any body of water.

1/22/71
8510 283


... AND
... FOR CONTROL
OF INSECTS AND DISEASES
OF VEGETABLES

CAUTION:
KEEP OUT OF REACH
OF CHILDREN

See additional cautions and warnings
on side panel.

ACTIVE INGREDIENTS:
*Malathion (O,O-dimethyl dithio-
phosphate of diethyl
mercaptosuccinate) 5.6%
**Methoxychlor, Technical 5.0%
Captan (N-trichloromethylthio-4-
cyclohexene-1,2-dicarboxi-
mide) 7.0%
INERT INGREDIENTS: 82.4%
100.0%
**Equivalent to 4.4% 2,2-bis(p-methoxy-
phenyl)-1,1,1-trichloroethane and 0.6%
other isomers and related compounds.

GARDEN VEGETABLE DUST



NET WEIGHT 1 POUND



DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Agway Vegetable Garden Dust Improved is for the control of vegetable insects and diseases as listed below. For best results, apply at 7-10 day intervals throughout the growing season. Cover the entire plant including the undersides of the leaves unless a soil application is indicated. Do not overtreat. Overtreating may cause injury to plants.

At time of harvest, remove excessive residue by washing or brushing. The following application rates may be measured by using 5 level tablespoons per ounce.

For control of Aphids, Bean Leaf Beetles, Leafhoppers, and Mexican Bean Beetles. Apply 8 ozs. per 1,000 sq. ft. when insects first appear and repeat as needed. Do not apply within 3 days of harvest. For the control of Damping-off fungi. Apply 21 ozs. per 1,000 sq. ft. of soil surface and work into the top 3-4 inches before planting.

For control of Flea Beetles. Apply 8 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 7 days of harvest.

For the control of Aphids. Apply 10 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 7 days of harvest.

For the control of Aphids and Flea Beetles. Apply 10½ ozs. per 1,000 sq. ft. when insects first appear. For the control of Corn Earworm, Fall Armyworm and Sap Beetles. Apply 10½ ozs. per 1,000 sq. ft. when 10% of the ears show early silk. Make 3 to 4 applications, applying at 2 day intervals. Do not apply within 10 days of harvest. Do not feed treated forage to dairy animals or to animals being fitted for slaughter.

For the control of Aphids, Cucumber Beetles, Leafhoppers, Squash Bug, Squash Vine Borers, Spider Mites, Angular Leaf Spot, Anthracnose, and Downy Mildew. Apply 14½ ozs. per 1,000 sq. ft. Make first application when initial leaves appear. Direct the dust toward the stems at the base of the plants and on the vines. The plants should be dry at the time of application. Repeat the treatment at 7-10 days throughout the growing season. Do not apply within 7 days of harvest.

For the control of Aphids, Flea Beetles, Leafhoppers, Pepper Maggots, Tomato Fruitworm, Tomato Hornworm, Spider Mites, Tomato Russet Mite, Tomato Early and Late Blight, Anthracnose and Septoria Leaf Spot. On Tomatoes, apply 16 ounces per 1,000 sq. ft. On Eggplant, apply 12 ounces per 1,000 sq. ft. On Peppers apply 9½ ounces per 1,000 sq. ft. Apply at first sign of insects damage or infection and repeat at 7 to 10 day intervals throughout the growing season. Do not apply within 7 days of harvest.

For control of Aphids, Cabbage Loopers, Leafhoppers, and Mites. Apply 10½ ozs. per 1,000 sq. ft. Repeat applications at 7 to 10 day intervals. Do not apply within 14 days of harvest.

For control of Aphids. Apply 12 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 14 days of harvest.

For control of Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight. Apply 16 ozs. per 1,000 sq. ft. Make first application when insects or infection appears and repeat at 7 to 10 day intervals throughout the growing season.

For control of Aphids, Lygus Bugs, Spittlebugs, Strawberry Leaf Rollers, Strawberry Weevils, Fruit Rot and Leaf Spot. Apply 13 ozs. per 1,000 sq. ft. when new growth appears and repeat at 7 to 10 day intervals as necessary for control. Do not apply within 14 days of harvest. At harvest time, remove excessive residue by washing fruit.