


AGWAY VEGETABLE GARDEN DUST

Contains Malathion, Methoxychlor* and Captan for Control of Insects and Diseases of Vegetables.

1081
PA12
8590-283



ACTIVE INGREDIENTS:

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	5.6%
*Methoxychlor, Technical	5.0%
Captan (N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide)	6.6%
Related Compounds	0.4%
INERT INGREDIENTS:	82.4%
TOTAL	100.0%

*Equivalent to 4.4% 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 0.6% other isomers and related compounds.

CAUTION
KEEP OUT OF REACH OF CHILDREN.
See additional cautions and warnings on side panel.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if inhaled, swallowed or absorbed through skin. Avoid skin contact with dust. Avoid breathing of dust. Wash thoroughly after using and before eating. Avoid contamination of feed and foodstuffs.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. Apply respiration, if indicated.

If on skin, remove contaminated clothing and wash affected areas with soap and water.

If in eyes, flush eyes for at least 15 minutes with water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Vegetable Garden Dust is for the control of vegetable insects and diseases as listed below. For best results, apply at 7-10 day intervals throughout the growing season. Cover the entire plant including the undersides of the leaves unless a soil application is indicated. Do not over-treat. Over-treating may cause injury to plants. At time of harvest, remove excessive residue by washing or brushing. The following application rates may be measured by using 5 level tablespoons per ounce.

Avoid use on wet foliage or in presence of high humidity or high temperature of several days duration.

BEANS: For control of Aphids, Bean Leaf Beetles, Leafhoppers, and Mexican Bean Beetles - Apply 8 oz. per 1,000 sq. ft. when insects first appear and repeat as needed. Do not apply within 3 days of harvest. For the control of Damping-off fungi - apply 21 oz. per 1,000 sq. ft. of soil surface and work into the top 3-4 inches before planting.

BEETS: For control of Flea Beetles. Apply 8 oz. per 1,000 sq. ft. when insects first appear. Do not apply within 7 days of harvest.

CELERY: For the control of Aphids. Apply 10 oz. per 1,000 sq. ft. when insects first appear. Do not apply within 8 days of harvest.

CORN: For the control of Aphids and Flea Beetles. Apply 10½ oz. per 1,000 sq. ft. when insects first appear. For the control of Corn Earworm, Fall Armyworm and Sap Beetles. Apply 10½ oz. per 1,000 sq. ft. when 10% of the ears show early silk. Make 3 to 4 applications, applying at 2 day intervals. Do not apply within 10 days of harvest. Do not feed treated forage to dairy animals or to animals being fitted for slaughter.

CUCUMBER, SUMMER SQUASH, PUMPKINS AND MELONS: For the control of Aphids, Cucumber Beetles, Leafhoppers, Squash Bug, Squash Vine Borers, Spider Mites, Angular Leaf Spot, Anthracnose, and Downy Mildew. Apply 14½ oz. per 1,000 sq. ft. Make first application when initial leaves appear. Direct the dust toward the stems at the base of the plants and on the vines. The plants should be dry at the time of application. Repeat the treatment at 7-10 days throughout the growing season. Do not apply within 7 days of harvest.

TOMATOES, PEPPERS, AND EGGPLANT: For the control of Aphids, Flea Beetles, Leafhoppers, Pepper Maggots, Tomato Fruitworm, Tomato Hornworm, Spider Mites, Tomato Russet Mite, Tomato Early and Late Blight, Anthracnose and Septoria Leaf Spot. On Tomatoes, apply 16 oz. per 1,000 sq. ft. On Eggplant, apply 12 oz. per 1,000 sq. ft. On Peppers, apply 9½ oz. per 1,000 sq. ft. Apply at first sign of insects damage or infection and repeat at 7 to 10 day intervals throughout the growing season. Do not apply within 7 days of harvest.

LETTUCE: For control of Aphids, Cabbage Loopers, Leafhoppers, and Mites. Apply 10½ oz. per 1,000 sq. ft. Repeat applications at 7 to 10 day intervals. Do not apply within 14 days of harvest.

SPINACH: For control of Aphids. Apply 12 oz. per 1,000 sq. ft. when insects first appear. Do not apply within 14 days of harvest.

POTATOES: For control of Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight. Apply 16 oz. per 1,000 sq. ft. Make first application when insects or infection appears and repeat at 7 to 10 day intervals throughout the growing season.

STRAWBERRIES: For control of Aphids, Lygus Bugs, Spittlebugs, Strawberry Leaf Rollers, Strawberry Weevils, Fruit Rot and Leaf Spot. Apply 13 oz. per 1,000 sq. ft. when new growth appears and repeat at 7 to 10 day intervals as necessary for control. Do not apply within 14 days of harvest. At harvest time, remove excessive residue by washing fruit.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-283

EPA Est. No. 8590-PA-1



AGWAY INC. — CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

NET WEIGHT 25 POUNDS

A 7/81