


# AGWAY

# Vegetable Garden Dust

Contains Malathion, Methoxychlor and Captan for control of insects and diseases of vegetables



KEEP OUT OF THE REACH OF CHILDREN.

**DANGER**

Causes irreversible eye damage

See additional precautionary statements on back panel

**Net Wt. 3 lbs.  
(1.36 kg)**

**ACTIVE INGREDIENTS**

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) .....	5.6%
*Methoxychlor, Technical .....	5.0%
Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide) .....	6.6%
Related Derivative .....	0.4%
<b>INERT INGREDIENTS:</b> .....	82.4%
	100.0%

\* Equivalent to 4.4% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related compounds

# AGWAY Vegetable Garden Dust

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

**DANGER:** Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin & clothing. Remove & separately launder clothing before reuse.

### STATEMENT OF PRACTICAL TREATMENT

If **swallowed**, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If **Inhaled**, remove victim to fresh air. Apply artificial respiration, if indicated.

If **on skin**, remove contaminated clothing and wash affected areas with soap and water.

If **in eyes**, flush eyes for at least 15 minutes with water. Call a physician immediately.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift & runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### ENDANGERED SPECIES RESTRICTIONS:

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

**Slackwater Darter**, Alabama (Lauderdale, Limestone and Madison), Tennessee (Lawrence and Wayne)

**Pecos gambusia**, New Mexico (Chaves and Eddy), Texas (Reeves, Jeff Davis and Pecos)

**Freshwater Mussels**, Tennessee (Hancock Claiborne, Hawkins and Sullivan)

Virginia (Smyth, Scott, Lee, Washington and Grayson)

**Comanche Springs Pupfish**, Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

**DISCLAIMER:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**NOTE:** This pesticide is to be sold **ONLY** in this original unbroken package.

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS & DIRECTIONS FOR USE.**

## STORAGE & DISPOSAL

Store in a cool, dry area preferably locked and out of the reach of children. Protect from excess heat.

Do not re-use empty container. Wrap container and put in trash collection.

Wear natural or synthetic rubber, vinyl or plastic gloves impermeable to Captan when using indoors or outdoors. When using outdoors wear long pants & long sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap & water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

Agway Vegetable Garden Dust is for the control of vegetable insects and diseases as listed below. For best results, apply at 7-10 day intervals throughout the growing season. Cover the entire plant including the undersides of the leaves unless a soil application is indicated. Do not over-treat. Overtreating may cause injury to plants. At time of harvest, remove excessive residue by washing or brushing. The following application rates may be measured by using 5 level tablespoons per ounce.

**BEANS:** For control of Aphids, Bean Leaf Beetles, Leafhoppers, and Mexican Bean Beetles. Apply 8 ozs. per 1,000 sq. ft. when insects first appear and repeat as needed. Do not apply within 3 days of harvest. For the control of Damping-off fungi. Apply 21 ozs. per 1,000 sq. ft. of soil surface and work into the top 3-4 inches before planting.

**BEETS:** For control of Flea Beetles. Apply 8 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 7 days of harvest.

**CELERY:** For the control of Aphids. Apply 10 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 7 days of harvest.

**CORN:** For the control of Aphids and Flea Beetles. Apply 10½ ozs. per 1,000 sq. ft. when insects first appear. For the control of Corn Earworm, Fall Armyworm and Sap Beetles. Apply 10½ ozs. per 1,000 sq. ft. when 10% of the ears show early silk. Make 3 to 4

applications, applying at 2 day intervals. Do not apply within 10 days of harvest. Do not feed treated forage to dairy animals or to animals being fitted for slaughter.

**CUCUMBERS, SUMMER SQUASH, PUMPKINS AND MELONS:** For the control of Aphids, Cucumber Beetles, Leafhoppers, Squash Bug, Squash Vine Borers, Spider Mites, Angular Leaf Spot, Anthracnose, and Downy Mildew. Apply 14½ ozs. per 1,000 sq. ft. Make first application when initial leaves appear. Direct the dust toward the stems at the base of the plants and on the vines. The plants should be dry at the time of application. Repeat the treatment at 7-10 days throughout the growing season. Do not apply within 7 days of harvest.

**TOMATOES, PEPPERS, AND EGG-PLANT:** For the control of Aphids, Flea Beetles, Leafhoppers, Pepper Maggots, Tomato Fruitworm, Tomato Hornworm, Spider Mites, Tomato Russet Mite, Tomato Early and Late Blight, Anthracnose and Septoria Leaf Spot. On Tomatoes, apply 16 ozs. per 1,000 sq. ft. On Eggplant, apply 12 ozs. per 1,000 sq. ft. On Peppers, apply 9½ ozs. per 1,000 sq. ft. Apply at first sign of insect damage or infection and repeat at 7 to 10 day intervals throughout the growing season. Do not apply within 7 days of harvest.

**LETTUCE:** For control of Aphids, Cabbage Loopers, Leafhoppers, and Mites. Apply 10½ ozs. per 1,000 sq. ft. Repeat applications at 7 to 10 day intervals. Do not apply within 14 days of harvest.

**SPINACH:** For control of Aphids. Apply 12 ozs. per 1,000 sq. ft. when insects first appear. Do not apply within 14 days of harvest.

**POTATOES:** For control of Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers, Early and Late Blight. Apply 16 ozs. per 1,000 sq. ft. Make first application when insects or infection appears and repeat at 7 to 10 day intervals throughout the growing season.

**STRAWBERRIES:** For control of Aphids, Lygus Bugs, Spittlebugs, Strawberry Leaf Rollers, Strawberry Weevils, Fruit Rot and Leaf Spot. Apply 13 ozs. per 1,000 sq. ft. when new growth appears and repeat at 7 to 10 day intervals as necessary for control. Do not apply within 14 days of harvest. At harvest time, remove excessive residue by washing fruit.



Agway Inc.  
PO Box 4741, Syracuse, NY 13221