

FOR: LABEL COPY - Thiram - Tinasad 5-5 G

DATE: 11/20/67

(((REAR PANEL)))

DANGER: (((18 pt. type))) May be fatal if swallowed, inhaled, or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for protection. Keep all unprotected persons out of the operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs.

This product is toxic to fish and wildlife. Do not contaminate any body of water. Birds feeding on treated areas may be harmed. Use this product only as specified on this label. (((Add paragraph at bottom of p.

4 as part of this paragraph)) (((Leave space before "First Aid")))

FIRST AID TREATMENT: SYMPTOMS: A sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea. TREATMENT: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, induce vomiting by giving a tablespoonful of salt in a glass of warm water. Administer milk or water freely and again induce vomiting until vomit fluid is clear. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash with flowing water for at least 15 minutes.

NOTE TO PHYSICIAN: ANTIDOTE: Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

Compound inhibits cholinesterase resulting in stimulation of the central nervous system, the parasympathetic nervous system, and the somatic motor nerves. Do not give morphine. Watch for pulmonary edema, which may develop in serious cases of poisoning even after 12 hours. At first sign of pulmonary edema, the patient

ACCEPTED
 12/16/68
 ON THE FEDERAL INSECTICIDE
 FUNGICIDE AND MITICIDE ACT
 FOR CONTROL OF PESTS REGISTERED
 UNDER FEDERAL LAW SUBJECT
 TO ATTACHED COMMENTS.

PCA - 50 CHARACTERS
 ELITE - 60 CHARACTERS

nr

O.K BY

STATEMENT OF ANALYSIS OF PLANT FOOD

Total Nitrogen	10.00%
Available Phosphoric Acid (P ₂ O ₅)	20.00%
Soluble Potash (K ₂ O)	20.00%



POISON

DANGER:

KEEP OUT OF REACH OF CHILDREN

DANGER: Poisoning may occur if this material is swallowed, inhaled, or absorbed through the skin. Do not breathe dust. Do not get in eyes or on skin. Wear protective clothing and gloves and wash the clothing and gloves directly after use. If Agway Ferti-Flow 10-20-20 With 0.00% Insecticide is used on non-food crops in an enclosure, wear a U.S. Department of Agriculture approved respirator mask. Do not contaminate feed or food. Wash thoroughly after application or handling. May be absorbed through the skin.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not reuse. Do not reuse empty containers. Dispose by burying with wastes or by burning. Stay away from smoke. Apply this product only as specified on this label.

SYMPTOMS OF POISONING: Tightness of the chest, sweating, headache, blurred vision, nausea, vomiting, and diarrhea.

TREATMENTS: Call physician immediately. Induce vomiting with a tablespoon of salt in a glass of warm water if the patient has swallowed the poison. Continue to induce vomiting until water administered to the patient becomes clear. If contact is with the skin, remove clothes and wash area with soap and warm water. For contact with eyes, wash with flowing water for 15 minutes.

TO PHYSICIAN: Atropine sulfate is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

ANTIDOTE: Administer atropine sulfate in large therapeutic doses. Repeat as necessary to point of tolerance.

PATENT LICENSE: The principal active insecticidal ingredient of this product is patented under U.S. Patent No. 2,759,010 of Chemagro Corporation, Kansas City, Missouri, and the purchase of this product includes a limited license under said patent to use or reformulate the product only in the manner provided by the recommended applications on the present label. The purchaser obligates itself to notify Chemagro Corporation of any proposed use or reformulation of such licensed product not provided by said recommended applications, and to obtain Chemagro's approval in writing before making any such use or reformulation of the licensed product. Any use, or reformulation intended for use, of the licensed product in the home and garden field is specifically prohibited by said license.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from breach of this warranty shall be limited to direct damages, and shall not include consequential or commercial damages such as loss of profits or value, etc.

CAUTION: When used on beans and cowpeas at the maximum recommended rates under adverse conditions, such as extremely cool and wet or extremely dry weather, this product may cause some delay in emergence, stunting of seedlings, or reduction of stand. Such damage may be pronounced in light, sandy soils. Plant injury may also occur or may be enhanced when certain pre-emergent herbicides are used in conjunction with this product.

VEGETABLES:

BEANS: For the control of Aphids, Leafhoppers, larvae of the Mexican Bean Beetle, Mites, and Thrips -- (1) On Snap or Green Lima Beans, apply 105-330 lbs. per acre banded on both sides of seed furrow at planting time only; (2) On Dry Beans, apply 105-330 lbs. at planting time or as a side-dressing on both sides of the row after plants are established. Do not apply within 60 days of harvest. Do not apply directly to seed and do not apply more than once per season to snap and green beans. See important statement under Directions for Use.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER: For the control of Aphids, Flea Beetles, and Leafhoppers on Cauliflower, Broccoli, and Brussels Sprouts as well as Mites and Thrips on Brussels Sprouts and to control Aphids and Flea Beetles on Cabbage, apply 105 lbs. per acre. Make the application in a band on each side of the seed furrow at planting or as a side-dressing after plants become established. Do not apply directly to the seed. For the control of Root Aphids on Cabbage apply 250 lbs. per acre. Apply by drilling or side-dressing. May be applied once per season on Broccoli and Cabbage and twice per season on Brussels Sprouts and Cauliflower. When applied two times per season, a minimum interval of 21 days must be observed between treatments. Do not apply within 14 days of harvest for Broccoli, 30 days of harvest for Brussels Sprouts, 42 days of harvest for Cabbage, or 40 days of harvest for Cauliflower.

POTATOES: For the control of Aphids, Leafhoppers, Potato Psyllid, early season reduction of Colorado Potato Beetles and Flea Beetles other than Tuber Flea Beetles, apply 330-500 lbs. per acre. Apply in the furrow or in a band on each side of the seed furrow at planting or as a side-dressing after plants are established. Use the higher rate on muck or organic soils. Do not apply within 75 days of harvest.

SPINACH: For the control of Aphids, Leafhoppers, Mites and Thrips, apply 165 lbs. per acre in a band on each side of the seed furrow at planting time only. Do not apply directly to seed.

TOMATOES: For early season reduction of Colorado Potato Beetle, apply 330 lbs. per acre. For the control of Aphids, Flea Beetles, Leafhoppers, Leaf Miners and Mites apply 105-330 lbs. per acre for one treatment per season or 105-330 lbs. per acre for two treatments per season. Where two applications per season are required, a minimum period of 21 days must be observed between treatments. Apply maximum dosage as a band on each side of the seed furrow at planting time only. A side-dress application after plants have become established is permissible where two applications are made at the lower rates. Do not apply within 30 days of harvest. See important statement under Directions for Use.

FIELD CROPS:

ALFALFA: For the control of Aphids and Mites apply 105 lbs. per acre as a broadcast treatment. Allow a minimum period of 26 days between applications.

BARLEY AND OATS: For the control of Aphids and Mites, apply 105 lbs. per acre by drilling at planting time or in a planting or post-emergence broadcast. Apply no more than two times per season at a minimum interval of 21 days between applications. Do not apply within 60 days of harvest. Do not graze or cut forage within 30 days of treatment.

CORN: For control of Northern Corn Rootworm, apply 105 lbs. per acre at planting time as starter fertilizer. Do not forage or cut treated corn for feed or ensilage within 40 days after application. Do not harvest for human purposes within 100 days of treatment.

WHEAT: For control of Hessian Fly apply 105 lbs. per acre in the fall by drilling or broadcast at planting time.

Manufactured for
AGWAY, INC.
Syracuse, New York

USDA REG. NO. 100-0-00000

0-00000

© DISTRIBUTION is a Registered Trademark of
Fadenfabrikken Eger A.G., Chemagro Corporation, New York, N.Y.