



# ORCHARD MOUSE BAIT-COATED

A wax coated cracked corn bait for control of meadow mice (*Microtus pennsylvanicus*) for use only in the orchards of New England, New York, New Jersey and areas of similar environmental conditions.

**ACTIVE INGREDIENT:**

Zinc Phosphide ..... 2%

**INERT INGREDIENTS:** ..... 98%

100%

**WARNING: KEEP OUT OF REACH OF CHILDREN.**

See additional warnings.

**WARNING:** May be fatal if swallowed. Do not breathe dust or fumes. Avoid contact with skin. Wear rubber gloves. Do not contaminate feed and foodstuffs. Wash after handling. Baits should be placed in areas inaccessible to children, pets, and domestic animals or in tamperproof bait boxes. Not for use or storage in or around the home.

Follow label recommendations carefully as to timing and mode of application to assure best possible control, to adequately protect wildlife, domestic fowl and animals.

This product is toxic to wildlife. Birds and other wildlife feeding in treated areas may be killed. When broadcasting bait, scatter evenly and do not apply to bare ground.

Do not reuse empty container. Destroy it by burying.

## DIRECTIONS

**READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.**

**BROADCAST BAITING:** Apply with a mechanical seeding device. Use 8 to 10 lbs. bait per acre. Scatter bait evenly on both sides of tree at the drip line of the branches. Do not apply to bare soil. Watch the weather and try to bait during periods of fair weather.

Apply immediately after harvest, before leaves fall, grass and weed growth mat down. The application can be repeated if necessary.

**NET WEIGHT 50 POUNDS**

**Hand Treating of Trails:** Later in the fall after rains or the first snowfall have matted down the cover or weeds, locate the trails or runways established by the rodents. Treat each quarter of the tree by carefully opening the weeds or grass over the runway and placing one teaspoon of bait in surface trails or at the mouth of holes leading to underground borrow systems. Do this for each quarter in several places at 1 to 2 foot intervals. Baiting should be conducted at the rate of 2 to 3 pounds per acre.

**Orchard Mouse Bait — Coated** should be applied immediately after harvest before leaves fall, grass and weed growth mat down, unless trail baiting by hand is to be practiced.

**AIRPLANE BAITING:** Apply 8 to 10 lbs. bait per acre. Do not apply over roadways. Other conditions apply as per broadcast baiting.

**NOTE:** This pesticide is to be sold ONLY in this original unbroken package.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

**AGWAY INC., FERTILIZER-CHEMICAL DIVISION, BOX 1333, SYRACUSE, NEW YORK 13201**

EPA Reg. No. 8590-304

ACCEPTED  
1-26-73  
8590-304