

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY SKEET-AWAY

DATE: 10/27/88

((FRONT PANEL))

Net Contents  
One Quart

AGWAY

SKEET-AWAY

For Mosquito Control

Contains 2 lbs. active ingredient Methoxychlor and 2 lbs. active ingredient Malathion per gallon.

ACTIVE INGREDIENTS:

Methoxychlor*, Technical. . . . .	23.8%
Malathion: 0,0 - dimethyl phosphorodithioate of diethyl mercaptosuccinate. . . . .	23.8%
Xylene Range Aromatic Solvent . . . . .	43.5%

INERT INGREDIENTS: . . . . .	8.9%
	<u>100.0%</u>

\*23.8% methoxychlor technical is equivalent to 20.8% of 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane; 3.0% other isomers and reaction products.

KEEP OUT OF THE REACH OF CHILDREN (12 pt.)

CAUTION (18 pt)

See additional precautionary statements on rear panel.

EPA Reg. No. 8590-310  
EPA Est. No. 8590-PA-1

Agway, Inc.  
PO Box 4741  
Syracuse, NY 13221

**ACCEPTED**

MAY - 3 1989

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 8590-310

OK BY

# ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY SKEET-AWAY

DATE: 10/27/88

((WEAR PANEL))

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Do not breathe spray mist. Do not get in eyes or on skin.

### STATEMENT OF PRACTICAL TREATMENT

If Swallowed, call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING, since vomiting may cause aspiration pneumonia. Gastric lavage is indicated if this product is taken internally.

If on Skin, remove contaminated clothing and wash affected areas with soap and water. Get Medical Attention if irritation persists. Wash contaminated clothing before reuse.

If in Eyes, flush eyes with plenty of water. Get Medical Attention if irritation persists.

If Inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

**NOTE TO PHYSICIAN:** Malathion may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to Atropine.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

### PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

Food utensils (teaspoons, tablespoons) should not be used for food purposes after use in measuring pesticides.

FOR: AGWAY SKEET-AWAY

DATE: 10/27/88

((REAR PANEL - CONT.))

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

STORAGE AND DISPOSAL (12 pt)

Store unused product in original container only, preferably in a locked storage area inaccessible to children.

Do not reuse empty container. Securely wrap empty container in several layers of newspaper and discard in trash.

Skeet-Away may be used as a residual spray for the control of adult mosquitoes or as a mosquito larvicide (1) applied as a pre-hatch winter treatment for the control of "spring brood" mosquito larvae, and (2) applied as a surface treatment to water breeding sites when larvae are present during the spring and summer.

ADULT MOSQUITOES

Residual Sprays: Apply 1 3/4 ounces (3 1/2 tablespoons) of Skeet-Away in enough water to cover 1,000 sq. ft. using a knapsack-type sprayer. Cover trunks and stems of shrubs and trees, as well as grassy vegetation, fences, out-buildings, and other structures near the ground. Sprays should be made to the point of runoff, but not excessive runoff. Repeat applications weekly. Do not spray foliage of Ferns, Hickory, Viburnum spp., Petunias, White pine, or Maples.

Larval Mosquitoes: Skeet-Away may be mixed with water for spray application or may be applied in undiluted form to the mosquito source. Apply only to known mosquito sites. When used in diluted form, mix in sufficient water to obtain even coverage.

Pre-hatch - Make a single application during the winter months at a rate of 3/4 ounces (1 1/2 tablespoons) per 1,000 sq. ft.

# ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY SKEET-AWAY

DATE: 10/27/88

((REAR PANEL - CONT.))

Larvicide - Make one or more applications to the source, as needed, for succeeding broods during the summer and spring. Apply at the rate of 3/4 ounces (1 1/2 tablespoons) per 1,000 sq. ft.

DISCLAIMER: Buyer assumes all risk of use, storage, or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

OK BY