


ACCEPTED
4-72
8590-324



METHOXYCHLOR 50W

WETTABLE POWDER FOR DAIRY AND BEEF CATTLE BUILDINGS AND CROPS

ACTIVE INGREDIENT:
Methoxychlor, Technical 50%
INERT INGREDIENTS: 50%
100%

*50% Methoxychlor, Technical is equivalent to 44% 2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroethane and 6% other isomers and reduction products.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Wash thoroughly after handling. This product is toxic to fish. Keep out of lakes, ponds and streams. Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

DIRECTIONS

READ THESE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND RESTRICTIONS.

FARM BUILDINGS For controlling stable flies, houseflies and fleas, use 1 lb. in 2 1/2 gallons of water. Apply 1 gallon of spray mixture to approximately 500 sq. ft. of surface—treating walls, floors, cracks, crevices, baseboards, pet bedding, stalls and pens. Such heavy deposits usually are effective for 3 to 6 weeks, unless removed by weathering or other means. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying. The use of any residual fly spray should be supplemented with proper sanitary measures, including spraying or removal of breeding sources such as manure in and around buildings. Do not apply in poultry houses.

BRAIN STORAGE BINS To clean up infestations in empty storage bins, use 1 lb. Methoxychlor 50W in 2 1/2 gallons of water applied at the rate of 1 gallon of spray per 500 sq. ft. before products are stored. This aids in control of cadelle, flat grain beetle, saw-toothed grain beetle, lesser grain borer, red flour beetle, long-headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle.

LIVESTOCK

Direct Dusting

Dairy Cattle—Lactating animals—For control of hornflies, use one tablespoonful (not to exceed 1/2 ounce) of Agway Methoxychlor 50W per animal. Sprinkle on poll, neck (top and sides), and back and upper sides; rub in lightly to move powder beneath the hair. Allow three weeks between applications.

Young Dairy Cattle over 1 mo. of age and non-lactating Mature Dairy Cattle—Dilute 1 part of Agway Methoxychlor 50W with 3 parts talc and shake to mix completely. Make a thorough application to the poll, neck, back and upper sides of the animal and work dust into the hair. Do not apply during lactation or within three weeks of freshening.

Beef Cattle, Goats and Sheep—For control of hornflies, lice and fleas. Make a thorough application to the animal. Work dust into hair or wool. Treatments are usually effective for about three weeks. Do not apply to lactating goats. Application on sheep following shearing gives temporary relief from sheep ticks (keds).

Swine—For control of hog lice dilute 1 part Agway Methoxychlor 50W with 3 parts talc, mix thoroughly and make a liberal application to animals and pens.

Direct Spray or Dips

Beef Cattle, non-lactating Dairy Cattle, Goats, Sheep and Swine—For control of Cattle lice, Hog lice and Sheep ticks, use 2 lbs. of Agway Methoxychlor 50W in 25 gals. of water. Thoroughly wet the animals. For Beef and Dairy cattle use 2 qts. of mix per head on large animals and proportionately less on smaller ones. Do not apply during lactation or within 2 weeks of freshening and do not dip calves under one month of age. Do not apply to lactating goats. For control of Tail louse of cattle, use 4 to 6 lbs. of Agway Methoxychlor 50W in 25 gals. of water and apply as a spray on the infested tail only.

AGRICULTURAL CROPS

Vegetables—For control of the insects listed below use 3 lbs. of Agway Methoxychlor 50W per 100 gals. of water and apply as a thorough spray to upper and lower surfaces of foliage and

fruit at the rate of 100 gals. per acre. Repeat treatment at first sign of insect infestation and repeat at 7-10 day intervals as infestation warrants.

Crop	Insects	Interval to Harvest
Corn	European corn borer, Fall armyworm, Japanese beetle, Tobacco budworm	14 days
Cucurbit	Cucumber beetle, Squash vine borer	1 day
Cucumber	Cucumber beetle	
Eggplant	Flan beetle, Leafhopper	
Kohlrabi	Flan beetle	
Pepper	Flan beetle, Rose chafar	
Potatoes	Cucumber beetle, Squash vine borer	
Squash	Cucumber beetle, Squash vine borer	
Tomato	Blister beetle, Fall armyworm, Flan beetle, Japanese beetle, Tobacco budworm	

Asparagus Asparagus beetle. If applied later than 3 days before harvest, remove residues by weeding or blanching.

Beans Bean leaf beetle, Corn earworm, All greenbean, Japanese beetle, Leafhopper, Rose chafar, Mexican bean beetle, Cowpea cicada

Black-eyed peas Cowpea cicada

Cabbages Cabbage worm, Flan beetle

Carrot/Flower Cabbage worm, Flan beetle

Corn Fall armyworm, Flan beetle, Japanese beetle, Rose chafar

Melons Cucumber beetle, Melon worm

Peas Pea weevil

Radish and Rutabaga Flan beetle

Beets Blister beetle, Flan beetle

Broccoli Cabbage worm, Flan beetle

Brussels sprouts Cabbage worm, Flan beetle

Carrot Leafhopper

Collards and Kale Flan beetle

Lettuce Leafhopper

Spinach and Turnip Flan beetle

Tree Fruits For control of the following insects, apply 2-3 lbs. of Agway Methoxychlor 50W per 100 gals. of water as a thorough cover spray when adults first appear. Repeat at 7-10 day intervals as long as adults are present. Do not apply more than 24 lbs. of Agway Methoxychlor 50W per acre.

Crop	Insects	Interval to Harvest
Apple	Apple maggots	7 days
Cherry	Cherry fruit flies, Cherry fruitworm	

Apply the above rate of Agway Methoxychlor 50W initially at petal fall for control of the following insects. Make 2 or more additional applications at 7-10 day intervals if infestation warrants.

Crop	Insects	Interval to Harvest
Apple	Codling moth, Japanese beetle, Leafhopper, Plum curculio	7 days
Cherry	Plum curculio	
Pear	Codling moth, Pear slugs	
Plum and Prune	Plum curculio	
Quince	Codling moth	
Apricot	Plum curculio	21 days
Avocado		
Cashew		

Small Fruit For control of the following insects apply 2-3 lbs. of Agway Methoxychlor 50W per acre in 100 gals. of water as a thorough cover spray when insects first appear. Repeat applications at 7-10 day intervals as infestation warrants.

Crop	Insects	Interval to Harvest
Blackberry	Flan beetle, Japanese beetle, Rose chafar, Leafhopper	14 days
Blueberry	Flan beetle, Japanese beetle, Rose chafar	
Strawberry	Flan beetle, Japanese beetle, Rose chafar	
Raspberries	Flan beetle, Japanese beetle, Rose chafar	
Cranberries	Cranberry leafhopper, Leafhopper	
Custard apple	Flan beetle, Japanese beetle, Rose chafar	
Guava	Flan beetle, Japanese beetle, Rose chafar	

Field and Forage Crops For control of the following insects apply 2-6 lbs. of Agway Methoxychlor 50W per acre in 70 to 40 gals. of water when insects first appear. Do not treat during bloom.

Crop	Insects	Do not harvest or graze within 7 days of application
Alfalfa	Alfalfa caterpillar, Armyworm, Fall armyworm, Flan beetle, Japanese beetle, Leafhopper, Lygus bugs, Spittlebug. (For Alfalfa weevil larval control apply when 20-50% of the leaf tips show damage.)	
Clover	Alfalfa caterpillar, Armyworm, Fall armyworm, Leafhopper, Lygus bug, Spittlebug, Aphis, weevil larvae.	
Cowpea	Cowpea cicada, Mexican bean beetle, Pea weevil.	
Soybeans	Mexican bean beetle, Soybean caterpillar, Velvet bean caterpillar.	

Mosquito Control For use as a pre-hatch winter treatment for control of "spring brood" mosquito larvae in New England States, make a single application of 2 lbs. of Agway Methoxychlor 50W during the winter months. Apply only to known mosquito breeding sites. Agway Methoxychlor 50W should be mixed with water for spray application.

NOTE: This product is toxic to fish. Keep out of lakes, streams or ponds. Shrimp and crab will be killed at application rate recommended. Do not apply where these are important resources. Do not apply to water drainage areas where runoff or flooding will contaminate ponds, lakes, or streams. Do not contaminate water by cleaning of equipment, or disposal of wastes.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

EPA Reg. No. 0990-324

MANUFACTURED BY AGWAY, INC., CHEMICAL DIVISION, BOX 1333, SYRACUSE, N. Y. 13201