

[8590-354]

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS

GENERAL:

To use remove protective cap and depress button being sure to aim spray opening at the location to be sprayed. Direct spray at insects wherever possible, and under eaves, into nests, cracks, holes or crevices wherever insects are noticed. Spray for about 10 seconds or until surface is moist from insecticide. To prevent entrance of insects into the house spray around points where outside plumbing, faucets, enter house and into any cracks or crevices in foundation as well as along sills and ledges.

FOR OUTDOOR USE:

Application should be made in late evening when insects are at rest. Hold can upright. Remove cap. Point valve opening away from face. With wind at back, aim at nests in trees, bushes, under eaves, chimneys, cracks or crevices.

FOR INDOOR USE:

Point valve opening away from face. Aim directly at nests under rafters. Do not spray in areas where foods and feedstuffs are exposed or handled. Do not remain in treated areas. Ventilate the area after treatment is completed.

Thoroughly saturate nest with spray, contacting as many insects as possible. To prevent insects from falling on skin and clothes, do not stand directly under nest. Best results can be obtained if nest is sprayed in late evening when most insects are at rest. If reinfestation occurs, repeat treatment.

Can also be used in attics, warehouses, homes, restaurants, and food processing plants where food and feedstuffs are not exposed.

Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using the product in these areas, apply only when the facility is not in operation.

FOR HOUSEFLIES, MOSQUITOES AND GNATS:

Close all doors and windows. Point nozzle upward, direct the spray mist to all parts of the room, especially windows and other light sources which attract insects. Fill the room with mist, then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed. Sweep up and destroy fallen insects. Repeat treatment as necessary.


FOR CRAWLING INSECTS: Roaches, earwigs, grain mites, sowbugs, spiders and centipedes; spray thoroughly into hiding places, cracks, crevices, moist areas, openings around pipes and sinks, under refrigerators, baseboards and storage areas. Spray directly on insects where possible. Repeat as necessary.

ANTS: Spray trails, nests and points of entry. Spray on ants where possible. Repeat as necessary.

DISCLAIMER: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

MANUFACTURED FOR AGWAY INC. FERTILIZER-CHEMICAL DIVISION BOX 1333, SYRACUSE, N.Y. 13201

EPA Reg. No. 8590-354
EPA Est. No.



NET WEIGHT
12-14 oz. 1 pound
or 1 pound 4 oz.

Hornet and Wasp Spray - SP

SBP-1382* Pressurized Spray Insecticide 0.25%

Agway Hornet and Wasp Spray will kill wasps, hornets and yellow jackets.

This formulation utilizes a synthetic pyrethroid, SPB-1382* in a pressurized spray formulation. Delivers a long spray pattern up to 10 feet - aids in reaching nests.

ACTIVE INGREDIENTS:

(5-Benzyl-3-furyl)methyl 2,2-dimethyl-3-(2-methylpropenyl) cyclopropanecarboxylate . . . 0.250%
Related Compounds 0.030

Petroleum distillate 29.375
INERT INGREDIENTS 70.345

100.000

*Cis trans isomers ratio: max. 30% cis and min. 70% trans.

U.S. Patent No. 912,079

SBP-1382* Registered trademark of CPC International, Inc.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of spray.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° F may cause bursting. Never throw container into fire or incinerator.

Apply this product only as specified on this label.

Avoid contamination of food and foodstuffs. Remove pets, birds, and cover fish aquariums before spraying. Do not use on food crops.