



residual fly spray 2e

for long residual fly control

Contains 25.2 BAYTEX® fenthion (as technical).
 Pounds per U.S. Gallon -- 2 lbs. Baytex®/gallon EC
 For Use By: Pest Control Operators, Livestock Producers, Commercial
 Nurserymen, Public Health Officials

ACTIVE INGREDIENTS

fenthion	
O, O-Dimethyl O-[4-(methylthio)-m-tolyl] phosphorothioate*	25.2%
Aromatic Petroleum Distillate	69.8%
INERT INGREDIENTS	5.0%

*U.S. Patent No. 3,042,703; Canadian Patent No. 631,472
 Baytex is a registered trademark in U.S. and Canadian Pat. Offs. by
 Farbenfabriken Bayer A.G. Chemagro Corporation Licensee.

WARNING KEEP OUT OF REACH OF CHILDREN

(See additional cautions and warnings elsewhere on Label.)

Comments DIRECTION
 READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Agway Residual Fly Spray 2E insecticide is recommended for use wherever flies are a problem. It provides long residual control and is equally effective against DDT resistant and non-resistant varieties. The spray can be applied to any surface including freshly whitewashed areas, without loss of biological effectiveness.

MIXING--The spray concentrate forms an emulsion when diluted with water. The resulting spray is suitable for use in hand or power-operated sprayers. To mix with water, pour required amount of Residual Fly Spray 2E concentrate into full amount of water and then agitate.

DOSAGE Use specified dosage of Agway Residual Fly Spray 2E insecticide in the amount of water necessary to give uniform coverage. The type of equipment used will determine the concentration required.

SPRAYING For residual localized fly control, applications should be made to surfaces of buildings, screen doors, window frames, around outdoor light fixtures, garages, and other surfaces where these insects congregate.

HOUSE FLIES--For residual fly control, mix 4 to 8 ozs. in 1 gallon of water for application per 500 square feet of time. If in localized areas, in patios, other outdoor structural surfaces, in buildings including dairy and beef barns and poultry houses. Do not apply directly to animals. Re-

sidual control lasts 7 to 8 weeks for the lower rate and even longer at the higher rate. Repeat as necessary. Do not apply as a space spray.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

May be fatal if swallowed, inhaled, or absorbed through the skin. Rapidly absorbed through the skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with soap and warm water after handling. Wash clothing with soap and hot water before reuse. Avoid contamination of feed, food, or food handling equipment, of food handling equipment, of food, contaminated, clean thoroughly before reuse. Keep out of reach of children. Should persons become ill, obtain prompt medical aid.

TO PHYSICIAN:

ANTIDOTE--Administer atropine sulfate in large therapeutic doses. Repeat as necessary to the point of tolerance. 2-PAM is also antidotal and may be administered in conjunction with atropine.

RESTRICTIONS

1. Do not apply when weather conditions favor drift of spray.
2. Do not use inside homes or in buildings where food is processed or stored.
3. Do not apply directly to animals.
4. Do not use on or allow spray treatment to drift onto water supplies. Do not contaminate feed, drinking water, or milk-handling equipment.
5. Do not apply sprays to flowers or other ornamental plants.

This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

CONTAINER DISPOSAL

Rinse equipment and containers and dispose of wastes by burying in non-crop land away from water supplies. Dispose of empty container by perforating and burying with wastes.

Do not use container in any connection with feed, food, and drinking water.

MANUFACTURED BY AGWAY INC.
 CHEMICAL DIVISION, BOX 1111
 SYRACUSE, NEW YORK 13211
 EPA REG. NO. 33-100-000