



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

JAN 26 1994

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Beverly Randall
UNIVERSAL COOPERATIVES INC.
AGENT FOR: AGWAY INC.
BOX 460
MINNEAPOLIS, MN 55440

Subject:

Label Amend:nent Submission of 08/04/93 in Response to PR Notice 93-7

EPA Reg. No. 8590-377

AGWAY DACTHAL 5G TURF HERBICIDE

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling AND
- WITHIN one year from date of this acceptance.

20yle.

Page 2

Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,

Im Tompkins, Deputy Chief Registration Support Branch Registration Division (7505W)

Attachment

(5% DACTHAL GRANULES):

Agreey Dacthal 5G Turf Herbicide is a 5% granular formula-tion for use as a presence application for control of brabgrass and other annual grasses and certain broadleaved weeds on mineral soils.

ight, is recommended for use on turf and ornamental plantings.

CTIVE INGREDIENT:
Dimethyl ester of tetrachloroterephthalic acid*....

Dacthale — Registered trademark of the Diamond Shamrock Corporation.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

ROO NOT take Internally, Avoid contact with eyes. Wash after standling or using.

DO NOT store near feed and foodstuffs, seed fertilizers or other pesticides in order to prevent contamination. Do not graze on treated areas or feed plant refuse to livestock. Store is in dry place.

DO NOT reuse empty container. Destroy by burning in a safe place. Stay away from smoke.

Personal Protective Equipment:

Applicators and other hardlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

BEST AVAILABLE COP

Follow manufacturer's instructions for cleaning /maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED

8590

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

SENERAL INFORMATION

MEAS THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE USE STRICTLY IN ACCORDANCE WITH LABEL CAU-

ANY APPLICATION OF DACTHAL SHOULD BE MADE PRIOR TO WEED SEED GERMINATION OR FOLLOWING A CULTI-VITOR.

Little the control of grasses and some broadleaved weed species, such as amooth and hairy crabgrass, green foxtall, bellow foxtall, lovegrass, wischgrass, carpetweed, common schickweed, lambequerters, pursiane and Florida pussley, Anyway Dacthal 5G Turf Herbicide is recommended at the grate of 180 pounds per acre, 4.0 pounds per 1,000 square feet, of 190 pounds for 250 square feet. A slightly higher rate top, application may be required for control of barnyardgrass, sinnual bluegrass; ground cherry, dodder, browntop panicum, goosgrass, Johnsongrass from seed, redroot pigweed, sessanis (S. macrocarpa) and nodding spurge.

Agrees Agreey Decthel 56 Turf Herbicide is used at the recom-impleded rate, up to 4 months weed control can be expected: Becoming grass, Johnsongrass established from rhizones, nut-ingress, gallinsoga, wild mustard, requeed, smartweed and eveloption have not been effectively controlled with Agway Cocthal 56 Turf Herbicide.

Spraiders should be checked to determine whether they are applying the correct amount of Agway Dacthal 5G Turf Herbicide. The application should be made to a uniform weed-free its soil surface within two to three days maximum following cultivation to prevent subsurface germinating weeds from a becoming established before the Agway Dacthal 5G Turf Herbicide is activated. Disturbing the soil surface following population will reduce effectiveness. Rates of application application will reduce effectiveness. Rates of application application be used by banding.

Dacthal 5G: Turf Herbicide application, sprinkler irrigation is should be used for best results.

The gray Dacthal 5G Turf Herbicide can also be applied by the shaker method. The ideal way of doing this is to measure the prescribed quantity of Agway Dacthal 5G Turf Schending over this area: 1 pound treats 250 square feet farea: 25 feet by 10 feet), 1 cupful treats 50 square feet.

RECOMMENDATION'S

TURF: Apply uniformly to establish turfgrass at the rate of 200 pounds per acre in the early spring before crabgrass and other annual grasses and broadleaved weeds germinate. This usually coincides with the time forsythia blossoms start to fall. One application, applied before weed germination in early spring usually provides season long control. Where spring seeding is necessary there should be a delay of approximately 60 days after the application. If reseeding is done before 60 days after application of Agway Dacthal 5G Turf Herbicide, the amount of grass sown should be increased by 50% to insure thrifty stands.

Early spring application of Agway Dacthal SG Turf Herbicide may be applied to newly seeded turf after they exhibit

greening of newly sprouted turigrass. Early spring applica-tion of Agway Dacthal 5G Turf Herbicide allows fall sowing of permanent grass. To control annual bluegrass and other erratic or late germinating grasses and weeds, a late sum-mer or early fall application of 300 pounds per acre of Agway Dacthal 5G Turf Herbicide should supplement the early spring application.

The use of DACTHAL on bentgrass greens should be restricted to portions of greens infested with crabgrass and bentgreens which are known to be tolerant to DACTHAL. Application to Cohansey or Toronto Bents is not recommended. Do not use on Dichondra turf. At the recommended rate of application, DACTHAL herbicide does not show herbicide injury or discoloration to established common lawn grasses. No more than the recommended rate should be used.

Spreader :	Setting for Normal Rate	Setting for 1/2 Normal Rate (crisscross)	Setting for 1½ Normal Rate
HH 20" Drop Spreader Model #1662	41/2 ·	31/2	51/4 -
HH Broadcast Spreader 0-2/11 5 ft. swath	3	.21/2	• 4
Cyclone Spreader Model #B1	3	21/2	¹ 3½
True Temper Broadcast Model D-2711	3	21/2	4
Lawn Beauty 24" Drop Model 2455	5		6
Scott's 24" Drop Model 75-3	41/2	31/2	5¥ 2
Scott's 18" Drop Model 35-3	4	274	5
Sears Craftsman 24" Drop Spreader Model #45219202	214	1¾	314
Greenfield 20" Drop Model #205	5	4	6 •

BEST AVAILABLE COPY

Ornamental Plantings: Apply at lining-out, late fall, or early spring at the rate of 180 lbs. Agway Dacthal 5G Turf Herbicide per acre (4 lbs. per 1,000 square feet). Applications should be made to soil recently cultivated to a uniform texture. These can be made immediately following lining-out of stock. With established plantings, application should be made following proper cultivation to remove existing weeds. Where possible, this should be done early in the spring. Late summer applications may prove beneficial for control of fall germinating weeds, if made following cultivation, Weed control up to 4 months or more may be expected following proper application.

Agway Decthal SG Turf Herbicide is specifically recommended for weed control with the following:

Abelia **Arborvitae** Ash Azalea Barberry Boxwood Birch Camellia Chestnut Cotoneaster Cottonwood Crab apple Cypress Deutzia Dogweed Eleeagnus Elm Euonymus, Fir Forsythia Gum Hawthorn Hydrangea Jap. andromeda Juniper Lantana Lilac

Locust Snapdragon Magnolia Strawflower Maple. Sunft ver Mock orange Verbe. Mountainlaurel Violet Oak Zinnia **Pachistima** Rhododendron Pine Rose Russian olive Pittosporum **Podocarous** Spirea **Poplar** Spruce Privet -Sycamore Redbud Tree peony Femleaf yarrow Tulip tree Forget-me-not Variegated privet Foxelove . Geranium Viburnum Walnut Weigela Willow Gold dust Golden tuft Lavendercotton Yew Baltic ivy Lupine Morningglory **Boston** ivy MOURNING bride English IVY Nasturtium Honeysuckle Peonies **Pachysandra** Pink heath Wilson ivy Purple coneflower Ageratum Salvia Alyssum

Aster
Babysbreath
Belifiower
Bleedingheart
Blood leaf
Bugloss
Candel larkspur
Candytuffs
Cinquefoil
Chrysanthemum
Columbine
Coleus
Coreopsis

Coral bells
Cosmos
Cuphea
Dahlia
Delphinium
Evening primrose
Feverlew
Four-O'clock
Gaillardia
Gladiolus
Gold, marguerite
Iris
Lily

Marigold Mother-of-Thyme Mourning pinks Orpine Petunia Poker plant Rose moss-Scarlet sage Stone Grop Sundrops Sweet Pea Virg. Spiderwort Wormwood

DO NOT USE on the following plants:

Bugle weed Button pink Carnation Geum

Germander Mesembryantheum Pansy Phlox Sweet William Telanthera

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages; and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

EPA Reg. No. 8590-377

MANUFACTURED BY AGWAY INC., CHEMICAL DIVISION, BOX 1333, SYRACUSE, NEW YORK 13201

NET WEIGHT FIFTY POUNDS

