

file



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

BEVERLY RANDALL  
UNIVERSAL COOPERATIVES, INC.  
AGENT FOR: AGWAY INC.  
BOX 460  
MINNEAPOLIS, MN. 55440

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 09/03/93 In Response to PR Notice 93-7  
EPA Reg. No. 8590-394  
DURSBAN 2EC

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

**WHAT THIS ACCEPTANCE MEANS:**

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

**WHAT YOU NEED TO DO NEXT:**

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable  
Printed with Soy-Cerata ink on paper that  
contains at least 50% recycled fiber

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (7505C)  
U.S. Environmental Protection Agency  
401 M Street, SW  
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Attachment

**AGWAY  
DURSBAN\* 2EC**

**Controls Pests of Households, Turf, Ornamental Plants as listed on this label  
-Also recommended for Mosquito Control and Area Control of Ticks.**

**To Be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts and Other Trained Personnel Responsible for Insect Control Programs.**

**Contains 2 pounds of chlorpyrifos per gallon.**

<b>ACTIVE INGREDIENT:</b>		
Chlorpyrifos [O,O-diethyl O-(3,5,6-trichloro-2-pyridyl)phosphorothioate]	.....	22.4%
<b>INERT INGREDIENTS**</b>	.....	<u>77.6%</u>
	<b>TOTAL</b>	<b>100.0%</b>

- \*\*Contains xylene range aromatic solvent
- \* DURSBAN® registered trademark of DowElanco
- U.S. Patent No. 3,244,586

**KEEP OUT OF REACH OF CHILDREN  
WARNING AVISO**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)**

**See Side Panel for Additional Precautionary Statements**

**EPA REG. NO. 8590-394  
EPA EST. 8590-PA-1**

**NET CONTENTS: 2 1/2 Gallons**

**AGWAY, INC.  
P.O. Box 4741  
Syracuse, NY 13221**

**ACCEPTED  
with COMMENTS  
in EPA Letter Dated  
JAN 28 1994**

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
8590-394**

48110

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS**

**WARNING: MAY BE ABSORBED THROUGH SKIN. MAY BE INJURIOUS TO EYES AND SKIN.**

**Do not take internally. Do not get in eyes or on skin. Flush contaminated eyes with plenty of water and get prompt medical attention. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Handle concentrate in a ventilated area.**

**Personal Protective Equipment:**

**Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category "G" on an EPA chemical resistance category selection chart.**

**Applicators and other handlers must wear:**

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as Barrier Laminate or Viton  $\geq 14$  mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- Chemical-resistant apron when cleaning equipment, mixing, or loading
- Dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C)

**Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.**

**User Safety Recommendations:**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**STATEMENT OF PRACTICAL TREATMENT  
ORGANOPHOSPHATE**

**If Swallowed:** Do not induce vomiting. Contains aromatic petroleum solvent. Call a physician immediately.

**If on Skin:** In case of contact, ~~remove contaminated clothing and~~ immediately wash skin with soap and water.

**If in Eyes:** Flush eyes with plenty of water. Call a physician.

**If Inhaled:** Remove victim to fresh air. Apply respiration if indicated.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

**PHYSICAL OR CHEMICAL HAZARDS  
COMBUSTIBLE**

**Do Not Use or Store Near Heat or Open Flame.**

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dursban 2EC is toxic to fish, birds and other wildlife. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Shrimp and crabs may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves, such as Barrier Laminate or Viton  $\geq$  14 mils
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

### **Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

- Keep children and pets off treated areas until spray has dried.
- Keep out of lawn and garden, fish pools and other bodies of water.

### PESTS INDOORS

General Information: Use Dursban 2EC insecticide as a spot treatment to control cockroaches, ants, crickets, firebrats, silverfish, spiders and brown dog ticks in and around residential, institutional and commercial buildings, including homes, apartment buildings, hospitals, stores and warehouses. Dilute to make a 0.25% or 0.5% chlorpyrifos spray as shown in the following SPRAY DILUTION CHART. Apply according to directions given for pests listed. Repeat the treatment as needed.

SPRAY DILUTION CHART

Gal of Spray Mixture Desired	Fluid Oz.* of Dursban 2EC to Use	
	0.25% spray	0.5% spray
Water-base spray:		
1	1 1/2	1 1/2
3	4	8
5	6 1/2	13 1/2
10	13 1/2	26 1/2
Oil-base spr.**:		
1	1	2
3	3	6
5	5	10
10	10	21

\*Note: 1 fluid ounce = 2 tablespoonfuls = 6 teaspoonfuls. Use only deodorized kerosene for oil based sprays to be applied in food handling establishments.

\*\* Calculated on basis of deodorized kerosene.

NOTE: Where quick "knockdown" and/or "flushing action" is desired, dichlorvos (DDVP) or pyrethrins (synergized pyrethrum) may be added to the spray mixture containing Dursban 2EC insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays; use pyrethrins at 0.25% in water-base sprays and 0.05% in oil-base sprays. Consult labeling for the dichlorvos or pyrethrins product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products.

Applications Within Food Handling Establishments (Places Other Than Private Residences In Which Food Is Held, Processed, Prepared or Served) Including, But Not Limited to Restaurants, Grocery Stores, Bakeries, Bottling Plants, Canneries and Grain Mills: Use as a spot treatment in and around both food and nonfood areas to control cockroaches (including strains resistant to certain other insecticides), ants, crickets, firebrats, silverfish, spiders and stored product insects such as confused flour beetles, sawtoothed grain beetles, Indian meal moths, rice weevils, Mediterranean flour moths and red flour beetles. Apply as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen or are suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. Do not introduce the spray into the air or allow the spray to contact food or food-contacting surfaces. Areas treated may include dark corners of rooms and closets; along and behind baseboards; beneath and behind stoves, refrigerator units and similar food preparation areas and equipment, floor drains; and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass crack and crevice treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and base or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling

establishments. In case of emergency (call back), Dursban 2EC may be applied after 2 days from last treatment but emergency use should be limited to once per month.

Applications of this product in food handling establishments other than as a spot and/or crack and crevice treatment are not permitted.

**Application Within Residential Buildings Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional, and Commercial Buildings Including Hospitals, Stores, Manufacturing Plants and Warehouses:**  
**For Spot Treatment Only** — Apply as a coarse spray or with a paint brush to localized areas where cockroaches (including strains resistant to certain other insecticides), ants, clover mites, crickets, firebrats, silverfish, and spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators and cabinets; around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For clover mites use a 0.5% spray only. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

For control of carpet beetles, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs, and furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

**Brown Dog Ticks:** Thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters; nearby cracks and crevices; along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

**ATTENTION (Pests Indoors):** Do not apply water base sprays of this product in conduits, motor housings, junction and switch boxes or other electrical equipment because of possible shock hazard. If there is need to treat such areas use only oil-base sprays. Prior to application deenergize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position. For Spot Application Only. Do not use in poultry houses, other animal buildings or in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

**PESTS OUTDOORS:**

**General Information:** Use Dursban 2EC for control of certain pests of lawns and ornamental plants, for area control of ticks and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

**Pests on Outside Surfaces of Buildings:** Use Dursban 2EC at the rate of 1 1/2 fluid ounces per one gallon of water (0.5% of chlorpyrifos emulsion) to control ants, clover mites, cockroaches, crickets, earwigs, fleas, flies, millipedes, mosquitoes, sowbugs, spiders and ticks by application as a residual spray or with a paint brush to outside surfaces of buildings including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen. Do not treat or allow the spray to contact plants (lawns, flowering plants, shrubs, trees, evergreens, etc.) because of possible injury. Repeat treatments as needed to maintain effectiveness.

**Pests of Lawns and other Ornamental and Recreational Turf Grass Areas:** Use Dursban 2EC insecticide to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and sod webworms (lawn moths). Mix Dursban 2EC with water according to directions given in the following table and apply as a coarse, low pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; retreat when needed.

**DOSAGE & DILUTIONS FOR USE ON TURF**

Area to be Treated	Amount of Dursban 2EC	Obviate with Water To Give
500 sq. ft.	1/4 fl. oz.	15 gallons
1,000 sq. ft.	1/2 fl. oz.	30 gallons
5,000 sq. ft.	7/8 fl. oz.	150 gallons
10,700 sq. ft.	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

**DOSAGES AND DIRECTIONS FOR USE ON TURF**

Pest	Dosage Dursban 2EC diluted in 20 gal. water per 1,000 sq. ft.	Remarks
Turfgrass (Hyperodes) Weevil	3 fl. oz.	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist in your area.
White grubs (including European chafar, southern masked chafar and Japanese beetle larvae)	3 to 6 fl. oz.	For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide in the underlying soil.

**DOSAGES AND DIRECTIONS FOR USE ON TURF**

Pest	Dosage Dursban 2EC per 1,000 sq. ft.	Remarks
Ants Armyworms Chiggers Chinch bugs	1 1/2 fl. oz.	Spray when pests first appear; retreat when needed. For sod webworms delay



Clover mite	watering or mowing
Crickets	the treated area for
Cutworms	12 to 24 hours after
Earwigs	treatment.
Fleas	
Grasshoppers	
Sad webworms	
Ticks	

Bluegrass	3 to 6 fl. oz.	Spray early in the
billbug		season when adults
		first appear. Retreat
		when needed.

• 1 1/2 fl. oz. per 1,000 sq. ft. = 7 1/2 fl. oz. per 5,000 sq. ft. = 1 pint per 10,700 sq. ft. = 2 quarts per acre.

**NOTE:** To help prevent infestation of buildings by ants, earwigs, crickets, and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

**Pests of Ornamentals:** Use Dursban 2EC to control bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, mites, spittlebugs, thrips (exposed) and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute Dursban 2EC with water according to directions given in the following table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage of infested plants. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals, if needed. **NOTE:** Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

**DOSAGE AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS**

Pests Controlled		Amount of Dursban 2EC in Water To Make		
		3 gals	10 gals	100 gals
Aphids	Mites	0.5	1.6	1 pint*
Bagworms	Orange-striped	fl. oz.	fl. oz.	
Eastern tent caterpillar	oakworm			
Fall webworm	Spittlebugs			
Grasshoppers	Whiteflies			
	Yellownecked caterpillar			
Cutworms	Mimosa	1.0	3.2	1 qt.**
Leafhoppers	webworm	fl. oz.	fl. oz.	
Mealybugs	Redhumped caterpillar			
	Thrips (exposed)			

\* Equivalent to 4 ounces of chlorpyrifos per 100 gallons  
 \*\* Equivalent to 8 ounces of chlorpyrifos per 100 gallons.

**Note:** Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

**Wood Boring Insects Infesting Shrubs and Trees** such as Dogwood Borer, Lilac Borer, Ash Borer, Bronze Birch Borer, and Rhododendron Borer: Apply Dursban 2EC to the trunks and lower limbs of

trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of two quarts of Dursban 2EC in 100 gallons of water and apply uniformly as a coarse, low-pressure spray.

**DOSAGES AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS**

Pests Controlled	Amount of Dursban 2EC in Water To Make		
	3 gals	10 gals	100 gals
Ash borer	2.0	6.4	2 quarts***
Bronze Birch borer	fl. oz.	fl. oz.	
Dogwood borer			
Lilac borer			
Rhododendron borer			
Scale crawlers			
Oak lecanium			
Pine needle			
Tea			
White peach			
Peach tree borer	5.8	19.2	6 quarts****
	fl. oz.	fl. oz.	

\*\*\* Equivalent to 1 pound of chlorpyrifos per 100 gallons.

\*\*\*\* Equivalent to 3 pounds of chlorpyrifos per 100 gallons.

**Peach Tree Borer:** Apply Dursban 2EC to flowering trees and shrubs of the Genus Prunus in water as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat your area.

**Area Control of Ticks and Chiggers:** Use Dursban 2EC to control ticks infesting non-crop land areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or possible public health problem. Apply Dursban 2EC in water at the rate of 1 pint per acre (equivalent to 1 fluid ounce per 2,700 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer or other suitable hand or power spray equipment. Treat low underbrush, grassy areas, weeds and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40 to 100 gallons per acre.

**Native Elm Bark Beetle: Prevention of Overwintering** to prevent the adult beetle from overwintering in uninfested trees and to reduce beetle populations in disease free areas, apply Dursban 2EC insecticide as a 0.5% mixture of chlorpyrifos in water to the bottom 9 feet of trunk. Wet the trunk thoroughly, but not to runoff. Care should be taken to apply the spray right to the base of the root flare. Applications can be made with either a back pack mist blower or a hydraulic pressure sprayer from spring through to the early fall. **Prevention of Branch and Twig Feeding on High Value Elm Trees:** To reduce the twig and branch feeding on uninfested trees deemed to be high in value apply a 0.5% chlorpyrifos water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that

will give thorough coverage to the tree crown.  
**Mosquitoes:** Apply Dursban 2EC to mosquito infested, non-crop areas according to the conditions and rates shown in the following table. Mix Dursban 2EC with either water or oil, such as kerosene or No. 2 fuel oil, using sufficient spray volume to give uniform coverage of area treated. Apply the spray using suitable application equipment such as hand and power ground sprayers, mist applicators, and aerial spray equipment.

### CONDITIONS AND RATES TO USE FOR MOSQUITO CONTROL

Condition of Area to be Treated	Amount to Use per Acre		Number Acres 1 Gal of Dursban 2EC Will Treat
	Dursban 2EC (fluid ounces)	Carbofuran (pounds)	
Light Control Light to medium vegetative cover	0.8 to 1.6	0.025 to 0.075	160 to 80
Medium to heavy vegetative cover	1.6 to 3.2*	0.025 to 0.05*	80 to 40
Adult Control Light to medium vegetative cover	1.6	0.025	80
Medium to heavy vegetative cover	3.2*	0.05*	40

\*The higher dosage will also give faster knockdown and longer residual control where there is little or no vegetative cover.

**Warning (Mosquito Control):** Dursban 2EC is toxic to shrimp, crabs and fish. Do not apply to water containing crustaceans and fish. Consult your State Fish and Game Department before using this product for mosquito control.

### STORAGE & DISPOSAL

**Prohibitions:** Do not contaminate or store near food, feed, fertilizers, seeds, insecticides, fungicides or other pesticides. Open dumping is prohibited. Do not reuse empty container.

**Storage:** Store unused product in original container only. Store in a cool, dry area out of the reach of children. Not for storage in or around the home. Protect from excess heat.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture & dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**NOTE:** This pesticide is to be sold ONLY in this original, unbroken package.

**DISCLAIMER:** Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

EPA Reg. No. 8590-394 EPA Est. No. 8590-PA-1

**RQ-1  
CHLORPYRIFOS  
MIXTURE  
ORM-A NA2783**