

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. May be injurious to the eyes and skin. Combustible.

Do not take internally. Do not get in eyes or on skin. Do not smoke while using. Wash thoroughly after handling. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of water and get medical attention. Avoid breathing spray mist or vapors. Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feedstuffs, and domestic water supplies. Keep container closed. **KEEP AWAY FROM HEAT AND OPEN FLAME.** Handle concentrate in a ventilated area. Keep children and pets off treated areas until spray has dried.

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse empty container. Rinse with water and discard in a safe place, preferably by burying in a location away from domestic water supplies. Bury unused chemical and spray mixture as well as rinse wastes in same manner as container.

NOTE TO PHYSICIAN: Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Use Dursban 2EC Lawn Insecticide to control chinch bugs, sod webworms (lawn moths), earwigs, cutworms, infesting home lawns and other ornamental and recreational turfgrass areas. Mix Dursban 2EC Lawn Insecticide with water according to directions given in the following table and apply as a coarse (large droplet) spray using a low pressure sprayer such as a knapsack type, hose attached applicator or other similar equipment. For best results, the lawn should be moist at time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when needed. One application usually will control chinch bugs for up to 8 weeks.

DOSAGE and DILUTIONS for USE on TURF

Area to be Treated	Amount* of Dursban 2EC Lawn Insecticide	Dilute with Water to Give
500 sq. ft.	3/4 fl. oz.	15 gallons
1,000 sq. ft.	1 1/2 fl. oz.	30 gallons
5,000 sq. ft.	7 1/2 fl. oz.	150 gallons
10,700 sq. ft.	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

*1 fluid ounce = 2 tablespoonfuls = 6 teaspoonfuls

To control certain White Grubs (larvae of European chafer and Japanese beetle): Use two to four times the amounts of Dursban 2EC shown in the turf table, i.e. 3 to 6 fl. oz. per 1000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July or August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

NOTE: To help prevent infestation of buildings by earwigs treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

MANUFACTURED FOR AGWAY INC.
Fertilizer-Chemical Division
Box 1333, Syracuse, New York 13201

EPA Reg. No. 8590-395

EPA Est. No. 8590-PA-1

BIG BOY

SEVIN* 5% DUST

ACCEPTED

AUG 1 1969

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER 08620-33 SUBJECT TO ATTACHED COMMENTS

50 LBS. NET WEIGHT
THIS PACKAGE CONTAINS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%

100%

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

CAUTION KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

DIRECTIONS

ALFALFA, CLOVERS, COWPEAS, PEANUTS, SOYBEANS: Blister Beetles, Mexican Bean Beetle-10 to 20 lbs. per acre. Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetles, Green Cloverworm, Japanese Beetle, Leafhoppers, Three-cornered Alfalfa Hopper, Thrips, Velvetbean Caterpillar-20 lbs. per acre. Armyworms, Corn Earworm, Stink Bugs, Webworms-20 to 30 lbs. per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

CORN: Corn Earworm (Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations), Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetle, Leafhoppers, Flea Beetles-30 to 40 lbs. per acre. For insects attacking silks and ears apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation.

TOMATO: Flea Beetles, Leafhoppers, Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer-50 to 60 lbs. per acre.

TOBACCO: (In plant beds-except Kentucky): Tobacco Flea Beetle-Apply 1 to 2 lbs. per 100 sq. yds. when plants appear. Repeat as needed at 10 to 14 day intervals and just before transplanting. (In Fields-All Areas)-Broadcast Applications - Hornworms, Tobacco Flea Beetles-Use 15 to 25 lbs. per acre depending on size plants. Use higher dosage on large plants and repeat as needed. Good coverage is essential. Heavier applications may cause plant injury. If late applications are necessary, treat immediately after priming. Do not apply within 5 days of priming or 15 days of cutting. Avoid getting wet with plant juices when cutting the crop.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

MANUFACTURED FOR
ESCAMBIA CHEMICAL CORPORATION
ALBANY, GEORGIA
USDA REG. NO. 8620 33