

AGWAY

LAWN INSECT CONTROL WITH DURSBAN

Kills chinch bugs, cutworms, earwigs, sod webworms (lawn moths) and certain white grubs infesting home lawns and other ornamental turf areas; borer insects in trees and shrubs.

Dursban® - registered trademark of DowElanco - U.S. Patent No. 3,244,586.

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 22.4%

INERT INGREDIENTS: \* 77.6%  
100.0%


Contains 2 pounds of chlorpyrifos per gallon.

\*Contains Xylene Range Aromatic Solvent

KEEP OUT OF REACH OF CHILDREN

WARNING

See additional cautions and warnings on side panel.




Net Contents

Manufactured for:  
AGWAY INC.  
P. O. BOX 4741  
SYRACUSE, NY 13221

EPA Reg. No. 8590-395

EPA Est. No. 1386-0H-1



**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING:** May be fatal if swallowed. May be absorbed through skin. May be injurious to eyes and skin.

Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feedstuffs, and domestic water supplies. Keep children and pets off treated areas until spray has dried.

**STATEMENT OF PRACTICAL TREATMENT**


If swallowed, contact a physician or poison control center immediately. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

If on skin, wash affected areas with soap and water.

If inhaled, remove victim to fresh air. Apply respiration, if indicated.

If in eyes, flush eyes with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.



ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water or wetlands (swamps, potholes, bogs, marshes). Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by disposal of equipment wash water.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.



STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Agway Lawn Insect Control With Dursban kills insects listed in the following tables which infest home lawns and other ornamental and recreational turfgrass areas. Mix with water according to directions given in the following table and apply as a coarse (large droplet) spray using a low pressure sprayer such as a knapsack type, hose attached applicator or other similar equipment. For best results, the lawn should be moist at time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when needed. One application usually will control chinch bugs for up to 8 weeks.

DOSAGES and DIRECTIONS for USE on TURF

Pest	Dosage Per 1000 Sq. Ft.	Remarks
Turfgrass (Hyperodes weevil)	3 fl. oz.	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist in your area.

<p><b>White grubs</b>          (Including          European          chafer,          southern          masked chafer          and          Japanese          beetle          larvae)</p>	<p>3 to 6          fl. oz.</p>	<p>For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide in the underlying soil.</p>
---	------------------------------------	--

---

<p><b>Bluegrass          billbug</b></p>	<p>3 to 6          fl. oz.</p>	<p>Spray early in the season when adults first appear. Re-treat when needed.</p>
--	------------------------------------	--

---

<p><b>Ants</b>  <b>Armyworms</b>  <b>Chiggers</b>  <b>Chinch bugs</b>  <b>Clover mite</b>  <b>Crickets</b>  <b>Cutworms</b></p>	<p>1½ fl. oz.</p>	<p>Spray when pests first appear; re-treat when needed. For sod <del>worms</del> <sup>worms</sup> delay watering or mowing the treated area for 12 to 24 hours after treatment.</p>
---	-------------------	---

697

- Earwigs
- Fleas
- Grasshoppers
- Sod webworms
- Ticks

**DOSAGE AND DILUTIONS**

<u>Area to be treated</u>	<u>Amount* of this product</u>	<u>Dilute with Water to give</u>
500 sq. ft.	3/4 fl. oz.	15 gallons
1,000 sq. ft.	1 1/2 fl. oz.	30 gallons
5,000 sq. ft.	7 1/2 fl. oz.	150 gallons
10,700 sq. ft.	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

\* 1 fluid ounce = 2 tablespoons = 6 teaspoons

**NOTE:** To help prevent infestation of buildings by earwigs, treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

Wood Boring Insects infesting Shrubs and Trees such as Dogwood Borer, Lilac Borer, Ash Borer, Bronze Birch Borer and Rhododendron Borer: Apply Lawn Insect Control with Dursban to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of two quarts of Lawn Insect Control with Dursban in 100 gallons of water and apply uniformly as a coarse, low pressure spray.

7 9 7

DOSAGES and DILUTIONS for USE on ORNAMENTAL PLANTS

Amount of LAWN INSECT CONTROL  
WITH DURSBAN in Water to Make

Pests Controlled	3 Gals.	10 Gals.	100 Gals.
Ash borer	2.0 fl. oz.	6.4 fl. oz.	2 qts.+++
Bronze Birch borer			
Dogwood borer			
Lilac borer			
Rhododendron borer			
Scale crawlers:			
Oak lecanium			
Pine needle			
Tea			
White peach			
Peach tree borer	5.8 fl. oz.	19.2 fl. oz.	6 qts.+++

+++ Equivalent to 1 pound of chlorpyrifos per 100 gallons

++++ Equivalent to 3 pounds of chlorpyrifos per 100 gallons

Peach Tree Borer. Apply Lawn Insect Control with Dursban to flowering trees and shrubs of the Genus Prunus in water as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray.

Thoroughly wet all bark areas from ground level to scaffold limbs. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat your area.

DISCLAIMER: Buyer assumes all risks of use, storage or handling, not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.