

P1119

8590 395

1 of 8

ACCEPTED

JUN - 4 1992

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8590-395

((FRONT PANEL))

NET CONTENTS
8 oz.

AGWAY

LAWN INSECT CONTROL WITH DURSBAN

Kills chinch bugs, cutworms, earwigs, sod moths) and certain white grubs infesting home lawns and other ornamental turf areas; borer insects in trees and shrubs.

ACTIVE INGREDIENT :

Chlorpyrifos 0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl)	
phosphorothioate	22.4%

INERT INGREDIENTS **

77.6%

100.0%

R

*Dursban - registered trademark of the Dow Chemical Company - U.S. Patent No. 3,244,586.

**Contains xylene range aromatic solvent.

Contains 2 pounds of chlorpyrifos per gallon.


KEEP OUT OF REACH OF CHILDREN (12 pt)

WARNING (18 pt)

See additional cautions and warnings on side panel

AGWAY INC.

P. O. BOX 4741
SYRACUSE, NY 13221



((REAR PANEL))

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through skin. May be injurious to eyes and skin.

Do not spray vegetable gardens or other plants used for food purposes. Keep away from food, feedstuffs, and domestic water supplies. Keep children and pets off treated areas until spray has dried.


STATEMENT OF PRACTICAL TREATMENT

If swallowed, Drink 1 or 2 glasses of water & induce vomiting by touching back of throat with finger.

If on skin, wash affected areas with soap & water

If in eyes, flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN Active ingredient is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.



ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER;
USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY
STATEMENTS AND DIRECTIONS.



STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Agway Lawn Insect Control With Dursban kills insects listed in the following tables which infest home lawns and other ornamental and recreational turfgrass areas. Mix with water according to directions given in the following table and apply as a coarse (large droplet) spray using a low pressure sprayer such as a knapsack type, hose attached applicator or other similar equipment. For best results, the lawn should be moist at time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; re-treat when needed. One application usually will control chinch bugs for up to 8 weeks.

DOSAGES and DIRECTIONS for USE on TURF

Pest	Dosage Per 1000 Sq Ft	Remarks
Turfgrass (Hyperodes weevil	3 fl. oz.	Spray suspected problem areas in mid-April and again in mid-May or as recommended by your



local Agricultural Extension Service specialist in your area.

<p>White grubs (Including European chafer, southern masked chafer and Japanese beetle larvae)</p>	<p>3 to 6 fl. oz.</p>	<p>For white grubs spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist for your area. Immediately after spraying, irrigate the treated area with ½ to 1 inch of water to wash the insecticide in the underlying soil.</p>
---	-----------------------	--

<p>Ants Armyworms Chiggers Chinch bugs Clover mite</p>	<p>1½ fl. oz.</p>	<p>Spray when pests first appear; re-treat when needed. For sod webworms delay watering or</p>
--	-------------------	--




Cricket		mowing the treated
Cutworms		area for 12 to 24
Earwigs		hours after treat-
Fleas		ment.
Grasshoppers		
Sod webworms		
Ticks		

Bluegrass	3 to 6	Spray early in the
billbug	fl. oz.	season when adults
		first appear. Re-
		treat when needed.

1½ fl. oz. per 1000 sq. ft. = 7½ fl. oz. per 5000 sq. ft. = 1 pint per 10,700 sq. ft. = 2 quarts per acre.

NOTE: To help prevent infestation of buildings by earwigs, treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.


Wood Boring Insects Infesting Shrubs and Trees



such as Dogwood Borer, Lilac Borer, Ash Borer, Bronze Birch Borer and Rhododendron Borer:
 Apply Lawn Insect & Borer Control with Dursban to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area. Mix at the rate of two quarts of Lawn Insect Control with Dursban in 100 gallons of water and apply uniformly as a course, low-pressure spray.

DOSAGES and DILUTIONS for USE on ORNAMENTAL PLANTS

Pests Controlled	Amount of LAWN INSECT CONTROL WITH DURSBAN in Water to Make		
	3 Gals.	10 Gals.	100 Gals.
Ash borer	2.0 fl oz	6.4 fl oz	2 qts+++
Bronze Birch borer			
Dogwood borer			
Lilac borer			
Rhododendron borer			
Scale crawlers:			
Oak lecanium			
Pine needle			
Tea			
White peach			



Peach tree borer	5.8 fl oz	19.2 fl oz	6 gts+++
------------------	-----------	------------	----------

+++ Equivalent to 1 pound of chlorpyrifos per
100 gallons

++++ Equivalent to 3 pounds of chlorpyrifos per
100 gallons

Peach Tree Borer. Apply Lawn Insect

Control with Dursban to flowering trees and shrubs
of the Genus Prunus in water as a trunk spray
before newly hatched larvae enter the trees.

Apply as a coarse, low-pressure spray. Thoroughly
wet all bark areas from ground level to scaffold
limbs. Consult your State Agricultural Experi-
ment Station or Extension Service specialist for
proper time to treat your area.

DISCLAIMER: Buyer assumes all risks of use,
storage or handling not in strict accordance with
directions given herewith.

NOTE: This pesticide is to be sold ONLY in this
original unbroken package.

