



THRAN 65W

30 POUNDS

AGRICULTURAL FUNGICIDE

ACTIVE INGREDIENT:	65%
Thiram (Tetramethylthiuram disulfide)	65%
INERT INGREDIENT:	35%
TOTAL	<u>100%</u>

KEEP OUT OF REACH OF CHILDREN

Do not reuse empty container. Destroy by burying with wastes or by burning. Stay away from smoke.

**READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE.
USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARN-
INGS AND DIRECTIONS.**

DIRECTIONS FOR USE AS A SPRAY. ALL RECOMMENDATIONS BASED ON 100 GALLONS OF SPRAY EXCEPT AS NOTED.

APPLES: Scab — 1½ to 2 pounds in prebloom, calyx, early cover sprays; and later cover sprays.

Rust — Use scab recommendations through first cover and include a special bloom spray of $\frac{3}{4}$ pound.

Other Diseases — Use scab recommendations as long as any one or more of the following diseases may appear — Apple Blotch, Bitter Rot, Black Pox, Black Rot (fruit and foliage), Botryosphaeria Spot, Brooks Spot, Fly Speck and Sooty Blotch.

CELERY: Early Blight (Cercospora), Late Blight (Septoria) and
Damping-Off.

Plant Beds — 1½ pound. Apply 15 to 25 gallons/1200 sq. ft. at 3-day intervals.

PEACHES: Brown Rot — For blossom blight, apply 1½ to 2 pounds at 3 to 4 day intervals during bloom, timed to keep newly opened blossoms covered. For fruit rot, apply at petal-fall and two weeks later. Make one or more additional applications at 7-day intervals beginning about one month before harvest, or earlier if rot is apparent on green fruit.

Rhizopus Rot — Apply as directed for fruit phase of brown rot.

Post — Apply at petal-fall and repeat 3 to 6 times at 10 to 14-day intervals or until disease is well under control.

Do not apply later than 7 days before harvest.

STRAWBERRY: Gray Mold (Botrytis).—As a spray, use 2 to 2½ lbs. per 100 gallons of water applying at rates up to 200 gallons per acre. As a dust use a mixture containing 7.5% to 10% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite applying at the rate of 40 lbs. per acre. Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results.

If applied within 3 days of picking, remove residues by washing.

TOMATOES: *Botrytis* Gray Mold and *Rhizoctonia* Fruit Rot — 1½ pounds when plants are set in the field and at 3 to 5-day intervals. If late blight, gray leaf spot or other foliage diseases are present, use combination of 1 pound Thiram 65W with 1½ pounds of either maneb or zineb wettable powder.

DIRECTIONS FOR USE AS A SEED AND SOIL TREATMENT FOR CONTROL OF DAMPING-OFF, SEED DECAY, SEEDLING BLIGHTS, ONION SMUT AND SWEET POTATO STEM ROT.

Apply at recommended dosages using the dry powder as received or mixed with water to form a slurry. Treated seed should be labeled, "Do not use treated seed for food, feed or oil."

Beans, Lima — 2.5 oz./100 pounds applied dry.

Beans, String — 1.5 oz./100 pounds applied as a slurry.

Beans, Soya — 1.5 oz./100 pounds applied dry or as a slurry.

Onions — As a dust furrow treatment — $1\frac{1}{2}$ pounds/acre mineral
soils, 14,500 linear feet of row or 3 pounds/acre on muck soils,
14,500 linear feet of row. *onion spray*

Soybeans — 1.5 oz./100 pounds applied dry or as a slurry.

Sweet Potatoes — 1.2 pounds/7½ gal. water applied as a preplanting root dip for ½ minute. Plant sprouts promptly after treatment.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

**AGWAY INC.
FERTILIZER-CHEMICAL DIVISION
BOX 1333, SYRACUSE, NEW YORK 13201**

EPA Reg. No. 8590-432

SP 174