

ADVERTISING AND SALES PROMOTION SERVICES

8590-432 PM 21 1068

COPY

FOR:

AGWAY THIRAM 65W

DATE: 4/15/85

((FRONT PANEL))

NET CONTENTS

5#/30#

AGWAY

THIRAM 65W

Fruit, Vegetable & Turf Fungicide

ACTIVE INGREDIENT:

Thiram (Tetramethylthiuram disulfide) 65%

INERT INGREDIENT:

35%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See additional precautionary statements on rear panel.

ACCEPTED

JUN 27 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 8590-432

AGWAY INC.

CHEMICAL DIV.

BOX 4933

SYRACUSE, NY 13221

EPA Reg. No. 8590-432

EPA Est. No. 8590-PA-1

OK BY

SHEET NO. 1-8

ADVERTISING AND SALES PROMOTION SERVICES

COPY

2068

FOR:

AGWAY THIRAM 65W

DATE: 4/15/85

(((REAR PANEL)))

PICA - NO CHARACTER
ELITE - NO CHARACTER

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes eye & skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap & water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, Call a Physician or Poison Control Center immediately. Drink 1 or 2 glasses of water & induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If on skin, remove contaminated clothing & wash affected areas with soap & water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

If in eyes, flush eyes with plenty of water. Get medical attention if irritation persists.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

During commercial exposure in seed treatment & loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the mine Safety & Health Administration (formerly the U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under provisions of 30 CFR, Part 11 Wash hands, arms & face thoroughly with soap & water after handling & before eating or smoking. Routinely shower or bathe after work:

OK BY

SHEET NO

2-8

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY THIRAM 65W

DATE: 4/15/85

Wash all clothing with soap & water before reusing.

Consumption of Alcoholic beverages increases the toxic effects of Thiram.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water or wetlands.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ALL RECOMMENDATIONS BASED ON 100 GALLONS OF SPRAY EXCEPT AS NOTED.

PRAISE DO CHARACTER
ELITE DO CHARACTER

OK BY

SHEET NO.

3-8

ADVERTISING AND SALES PROMOTION SERVICES

4068
COPY

FOR: AGWAY THIRAM 65W

DATE: 4/15/85

RE-ENTRY

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

CAUTION

(Insert area or field description) treated with Thiram on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Thiram and describe his condition. For further information see the "Statement of Practical Treatment." Oral warnings must include the same information that is in the written warnings.

PICA 80-CHARACTER
ELECTRONIC CHARACTER

O.K. BY

SHEET NO.

FOR: AGWAY THIRAM 65W

DATE: 4/15/85

DIRECTIONS FOR USE AS A SPRAY ON FRUITS & VEGE-
TABLES:

APPLES:

Scab - 1½ to 2 pounds in pre-bloom, calyx, early
cover sprays; and later cover sprays.

O.K. BY

Rust - Use scab recommendations through first
cover and include a special bloom spray of ¾
pound.

Other Diseases - Use scab recommendations as
long as any one or more of the following diseases
may appear - Apple Blotch, Bitter Rot, Black Pox,
Black Rot (fruit and Foliage), Botryosphaeria
Spot, Brooks Spot, Fly Speck and Sooty Blotch.

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

CELERY: Early Blight (Cercospora), Late Blight
(Septoria) and Damping-Off.

Plant Beds - 1½ pounds. Apply 15 to 25 gallons/
1200 sq. ft. at 3-day intervals.

PEACHES:

Brown Rot - For blossom blight, apply 1½ to 2
pounds at 3 to 4 day intervals during bloom,
timed to keep newly opened blossoms covered. For
fruit rot, apply at petal-fall and two weeks later.
Make one or more additional applications at 7-day
intervals beginning about one month before har-
vest, or earlier if rot is apparent on green fruit.

Rhizopus Rot - Apply as directed for fruit phase
of brown rot.

O.K. BY

SHEET NO.

5-8

ADVERTISING AND SALES PROMOTION SERVICES

COPY ⁶⁴⁸

FOR: AGWAY THIRAM 65W

DATE: 4/15/85

Scab - Apply at petal-fall and repeat 3 to 6 times at 10 to 14-day intervals or until disease is well under control.

Do not apply later than 7 days before harvest.

STRAWBERRIES:

Gray Mold (Botrytis) - As a spray, use 2 to 2½ lbs. per 100 gallons of water applying at rates up to 200 gallons per acre. As a dust use a mixture containing 7.5% to 10% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite applying at the rate of 40 lbs. per acre. Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results.

If applied within 3 days of picking, remove residues by washing.

TOMATOES:

Botrytis Gray Mold and Rhizoctonia Fruit Rot - 1½ pounds when plants are set in the field and at 3 to 5-day intervals. If late blight, gray leaf spot or other foliage diseases are present, use combination of 1 pound Thiram 65W with 1½ pounds of either maneb 80% or zineb 75% wettable powder.

Do not rotate treated crop with crops other than those with registered Thiram uses.

Do not reenter treated areas within 24 hours following application unless protective clothing is worn.

OK BY

SHEET NO.

ADVERTISING AND SALES PROMOTION SERVICES

COPY 7 of 8

FOR:

AGWAY THIRAM 65W

DATE: 4/15/85

DIRECTIONS FOR USE AS A SEED AND SOIL TREATMENT
FOR CONTROL OF DAMPING-OFF, SEED DECAY, SEEDLING
BLIGHTS, ONION SMUT AND SWEET POTATO STEM ROT.

Apply at recommended dosages using the dry powder
as received or mixed with water to form a slurry.
Treated seed should be labeled, "Do not use
treated seed for food, feed or oil."

Bears, Lima - 2.5 oz./100 pounds applied dry.

Beans, String - 1.5 oz./100 pounds applied as a
slurry.

Beans, Snap - 1.5 oz./100 pounds applied dry or
as a slurry.

Onions - As a dust furrow treatment - 1½ pounds/
acre mineral soils, (14,500 linear feet of row)
or 3 pounds/acre on muck soils (14,500 linear
feet of row.)

Soybeans - 1.5 oz./100 pounds applied dry or as
a slurry.

Sweet Potatoes - 1.2 pounds/7½ gal. water applied
as a preplanting root dip for ½ minute. Plant
sprouts promptly after treatment.

PICA - 80 CHARACTERS
ELITE - 80 CHARACTERS

OK BY

SHEET NO. 7-8

ADVERTISING AND SALES PROMOTION SERVICES

COPY 8 of 8

FOR: AGWAY THIRAM 65W

DATE: 4/15/85

STORAGE & DISPOSAL (12 pt)

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Store unused product in original container only. Store in a cool, dry area preferably locked & out of the reach of children and animals. Not for use or storage in or around the home. Protect from excess heat.

PESTICIDE DISPOSAL: Pesticide Wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

OK BY

SHEET NO.