

GRANULAR TURF INSECTICIDE DIAZINON® 5G

Controls: Chinch bugs, Sod Webworms, Lawn moths, Earwigs, White grubs of Japanese beetles, European chafers, Southern chafers, Billbugs and others as specified in label directions.

Ready-to-use. Contains 5% Diazinon insecticide.

ACTIVE INGREDIENTS:

0,0-diethyl 0-2-isopropyl-6-methyl-4-pyrimidinyl phosphorothioate* 5%

INERT INGREDIENTS

95%
100%

*Diazinon™ trademark of CIBA-Geigy Corp.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

OVERALL LAWN APPLICATION:

Agway Diazinon 5G controls the following lawn insects on both grass and dichondra lawns:

Chinch bugs, Earwigs, Ticks, Ants, Lawn moths, Sod webworms, Bermuda mites, Crickets, Armyworms, Brown dog ticks, Chiggers, Clover mites, cutworms, Fleas, Leafhoppers, Sowbugs, Millipedes, Springtails, Collembola.

Apply at the rate of 2 lbs. per 1000 sq. ft. (an area 20' by 50') or 3.2 oz. per 100 sq. ft. (an area 10' by 10'). For heavy lawns such as St. Augustine lawns, application should be increased to 3 lbs. per 1000 sq. ft. or 4.8 oz. per 100 sq. ft.

Hyperodes Weevil (New York Only)

Apply 2 lbs. per 1000 sq. ft. of grass. Water grass thoroughly after application. Treat problem areas in mid-April and again in mid-May.

Apply Agway Diazinon 5G to your lawn with a fertilizer spreader. Application should be made when grass is dry. Water lawn lightly after application. See rate chart below.

Method:

To help obtain uniform distribution, apply one-half the required amount of product per determined area to be

treated in one direction (refer to rate chart); then apply the balance at a right angle. Immediately after the application sprinkle the turf to wash the granules into the zone of insect infestation. Refer to table for gallons of water needed per given area. Repeat as necessary.

RATES TO USE

Area to be Treated	Pounds of Diazinon 5G	Gal. of water to sprinkle on turf after treatment
1,000 sq. ft.	2	30
5,000 sq. ft.	10	150
10,700 sq. ft.	20	325

SPREADER SETTING CHART

Broadcast (Rotaries)	Drop Types	
	once over	once over
Cyclone B	3.25	Agway 72" 5.00
MTD	3.25	MTD 5.00
Agway	3.25	Sears 19" 5.00
Central = 1675	3.25	Jackson 18" 5.00
		Agway 18" before 71 5.00
		Scott 5.00
		Central = 1622A 5.25
		Tru Temper 5.00

Billbugs and White Grubs of Japanese Beetles, European Chafer, Southern Chafer.

Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

RATES TO USE

Area to be Treated	Pounds of Diazinon 5G	Gal. of water to sprinkle on turf after treatment
1,000 sq. ft.	2.5	30
5,000 sq. ft.	12.5	150
10,700 sq. ft.	25	325

SPREADER SETTING CHART

Broadcast (Rotaries)	Drop Types	
	once over	once over
Cyclone B	3.75	Agway 72" 6.00
MTD	3.75	MTD 6.00
Agway	3.75	Sears 19" 5.75
Central = 1675	3.75	Jackson 18" 6.00
		Agway 18" before 72 6.00
		Scott 6.00
		Central = 1622A 6.25
		Tru Temper 6.00

SPOT TREATMENT

ANTS: Apply 1½ teaspoons of Agway Diazinon 5G over and around each individual ant hill.

DIGGER WASPS: Apply 1½ teaspoons of Agway Diazinon 5G in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants	Crickets	Millipedes
Brown Dog Ticks	Earwigs	Sowbugs
Clover Mites	Fleas	Springtails, Collembola

For control of the above insects, apply a five-foot band of Agway Diazinon 5G next to the foundation of the house. Distribute the granules uniformly, using a rate of 3.2 oz. for each 5' by 20' area to be treated.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under

abnormal conditions, or under conditions not foreseeable to seller and buyer assumes use. Any damages arising from a breach shall be limited to direct damages, and consequential commercial damages such as lost profits or values, etc.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination. Do not store near food or feed products. Avoid contact with skin, eyes, nose, or mouth. Wash thoroughly after handling. May be absorbed through skin. Keep children and pets off treated area until grass is washed into the soil and grass is dry.

Do not reuse container. Destroy it by incineration or by burning. Stay away from smoke.

Do not contaminate water by cleaning or disposal of wastes.

This product is toxic to fish and wildlife in lakes, streams or ponds. Birds feeding in treated areas may be killed.

Apply this product only as specified on label. Diazinon kills: Ants, Armyworms, Bermuda grass mites, Brown dog ticks, Chiggers, Chinch bugs, Crickets, Cutworms, Digger wasps, Earwigs, Fleas, Grasshoppers, Leafhoppers, Millipedes, Springtails, Sowbugs, Ticks, White grubs of: Japanese beetles, Southern chafers.

NOTE: This pesticide is to be sold only in its original unbroken package.

EPA Reg. No. 8590-443

AGWAY INC. FERTILIZER — CHEMICAL DIVISION
BOX 1333, SYRACUSE, N.Y. 13201

LAWN TURF INSECTICIDE DIAZINON® 5G

treated in one direction (refer to rate chart); then apply the balance at a right angle. Immediately after the application sprinkle the turf to wash the granules into the zone of insect infestation. Refer to table for gallons of water needed per given area. Repeat as necessary.

RATES TO USE

Area to be Treated	Pounds of Diazinon 5G	Gal. of water to sprinkle on turf after treatment
1,000 sq. ft.	2	30
5,000 sq. ft.	10	150
10,700 sq. ft.	20	325

SPREADER SETTING CHART

Broadcast (Rotaries)	Drop Types	once over	once over
Cyclone B	Agway (72)	3.25	5.00
MTD	MTD	3.25	5.00
Agway	Sears 19"	3.25	5.00
Central = 1675	Jackson 18"	3.25	5.00
	Agway 18"		5.00
	(before 71)		5.00
	Scott		5.00
	Central = 1622A		5.25
	Tru Temper		5.00

Billbugs and White Grubs of Japanese Beetles, European Chafer, Southern Chafer.

Apply 2.5 lbs. per 1000 sq. ft. of lawn. Water grass thoroughly after application. For grub control make application any time between late July and early October. For billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. Repeat as necessary.

RATES TO USE

Area to be Treated	Pounds of Diazinon 5G	Gal. of water to sprinkle on turf after treatment
1,000 sq. ft.	2½	30
5,000 sq. ft.	12½	150
10,700 sq. ft.	25	325

SPREADER SETTING CHART

Broadcast (Rotaries)	Drop Types	once over	once over
Cyclone B	Agway (72)	3.75	6.00
MTD	MTD	3.75	6.00
Agway	Sears 19"	3.75	5.75
Central = 1675	Jackson 18"	3.75	6.00
	Agway 18"		6.00
	(before 72)		6.00
	Scott		6.00
	Central = 1622A		6.25
	Tru Temper		6.00

SPOT TREATMENT

ANTS: Apply 1½ teaspoons of Agway Diazinon 5G over and around each individual ant hill.

DIGGER WASPS: Apply 1½ teaspoons of Agway Diazinon 5G in and around each individual ground nest opening. Make the application about dusk when the wasps are not active.

BAND TREATMENT AROUND FOUNDATION

The following insects are commonly found around or near the foundation of houses as well as in the lawn. Certain of these insects may even enter the house.

Ants	Crickets	Millipedes
Brown Dog Ticks	Earwigs	Sowbugs
Clover Mites	Fleas	Springtails (Collembola)

For control of the above insects, apply a five-foot band of Agway Diazinon 5G next to the foundation of the house. Distribute the granules uniformly using a rate of 3.2 oz. for each 5' by 20' area to be treated.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under

abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of food and feed. Do not store near food or feed products. Avoid inhalation of dust. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. May be absorbed through skin. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Do not reuse container. Destroy it by burying with wastes or by burning. Stay away from smoke.

Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed.

Apply this product only as specified on this label.

Diazinon kills: Ants, Armyworms, Bermuda mites, Billbugs, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Digger wasps, Earwigs, Fleas, Lawn moths, Leafhoppers, Millipedes, Springtails, Sod webworms, Sowbugs, Ticks, White grubs of: Japanese beetles, European chafers, Southern chafers.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590 413

EPA Est. No. 8590 PA 1

AGWAY INC.
FERTILIZER — CHEMICAL DIVISION
BOX 1333, SYRACUSE, N. Y. 13201