ACCEPTED

RECEIVED
29 OCT 1974
REGISTRATION DIVISION
PESTICIDES, EPA

AGWAY

DIAZINON® 10G

ACTIVE INGREDIENT
O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 10%
INERT INGREDIENTS 90%
..... 100%

®Trademark of CIBA Geigy Corp.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS, AND DIRECTIONS.

FRUITS

VEGETABLE FOLIAR INSECTS

Tomatoes: For control of ... apply 5-10 lbs per acre broadcast when ... Repeat at 7-14 day intervals or as required.

VEGETABLE SOIL INSECTS

... For control of cutworms in broccoli, cabbage, ... cucumbers, endive (escarole), kale, lettuce, ... peppers, snap beans, spinach, summer squash, ... watermelons, winter squash apply 20-30 lbs per ... broadcast just before planting. Work into ... for surface cutworms, 3-6 inches for subterranean cutworms.

... crickets: For control of mole crickets in broccoli, cabbage, cauliflower, collards, ... mustard greens, parsley, radishes, tomatoes, turnips apply ... 1-2 days before sowing seed or train planting ... immediately.

... maggots: For control of maggots in radishes apply 15-20 lbs per acre in ... furrow at planting.

... worms: For control of wireworms in broccoli, cabbage, cantaloupes, carrots, cauliflower, ... lettuce, melons, mustard greens, parsley, peppers, radishes, sweet ... watermelons, winter squash apply 20-30 lbs per ... broadcast at planting. Work into ... at planting for ... Band over row

FIELD AND FORAGE INSECTS

... and Corn: For control of ... surface and subterranean ... apply 20-30 ... at planting

... Apply ... 7 days ... to feed to ... immediately

... apply 1.25-2.5 oz per 10 ... at planting

... in Northern Corn Belt ... Apply ... for surface cut- ... 1-2 ... per ... before planting ... inches ... apply ...

... Apply 20 lbs per acre ... Do not ... 7 days ... to feed to ... immediately

Southwestern corn borer: Apply ... lbs per acre ... granules fall into whorls when second generation ... first adults ... one week later.

Lesser cornstalk borers: Apply 20-30 lbs per ... in a 10 inch band centered over row at planting or when ... Incorporate ... treated band with 1-2 inches of soil.

Corn may be harvested immediately following last application. Do not feed treated forage to dairy or beef cattle or sheep within 21 days after last application.

Cowpeas: For control of cutworms ... and subterranean ... apply 20-30 lbs per acre just before planting. Work into ... immediately, 2-3 inches for surface cutworms, 6 inches for subterranean cutworms.

Lespedeza: For control of cutworms ... and subterranean ... apply 20-30 lbs per acre broadcast just before planting and work into ... immediately, 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Peanuts: For control of southern corn rootworm ... apply 3 lbs per acre broadcast or ... over fruiting zone just before pegging. Work into top few inches of soil. Do not apply within 21 days of harvest.

... cornstalk borers apply 20-30 lbs per acre in a band over fruiting zone just before pegging. Cover band with 1-2 inches of soil. Do not apply within 21 days of harvest.

... chicks may be fed peanut hulls ... may be fed ... treated 7 days following treatment and ... may be fed peanut ... treatment.

Sorghum: For control of cutworms ... and subterranean ... apply 20-30 lbs per acre broadcast just before planting. Work into ... immediately, 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

... Apply 10-20 lbs per acre in a 10 inch band over row at planting or when ... emerges. Cover band with 1-2 inches of soil.

Do not ... apply within 7 days ... harvest. Forage may be fed to livestock immediately.

Soybeans: For control of ... apply 20-30 lbs per acre broadcast just before planting or when ... 1-2 inches of soil.

CAUTION KEEP OUT OF REACH OF CHILDREN

... Do not ... and ... Avoid ... Do not ... and clothes ... Wash hands ... after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

... Store in ... and keep ... Store ... away from ... and ... Do not ... Apply this ... only as ... on the label.

Do not reuse empty container. Destroy by burning with wastes or by burning. Stay away from smoke.

WARRANTY: ... warrants that the ... conforms ... description and ... reasonably ... the purposes ... in accordance with ... Agway ... nor any other ... assumes ... liability ... use of this ... contrary to ... or ... Buyer assumes ... risks ... such use. ... Agway's ... liability ... shall be limited ... the purchase price ... product involved.

NOTE: ... to be sold ONLY in ... original ... package.

EPA Reg. No. 8590-447 ... EPA Est. No. 8590-1

AGWAY INC.
FERTILIZER-CHEMICAL DIVISION, BOX 1333
SYRACUSE, NEW YORK 13201

NET WEIGHT 50 POUNDS