

# KELTHANE\* 1.5EC

**AGRICULTURAL MITICIDE CONTROLS MITES ON VEGETABLES, FRUITS, SHADE TREES, ORNAMENTALS, FLOWERS, AND CONTROLS CLOVER MITES IN HOMES.**

**ACTIVE INGREDIENTS:**

1, 1-bis (p-chlorophenyl)-2, 2, 2-trichloroethanol .....	18.5%
Xylene .....	73.0%
<b>INERT INGREDIENTS:</b> .....	8.5%
	<hr/> 100.0%

\*Kelthane, registered trademark of Rohm & Haas Co.

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

Use two teaspoons of Kelthane 1.5EC in one gallon of water unless otherwise noted. Spray both upper and lower plant surfaces when mites first appear. Repeat when necessary.

### FLOWERS, NURSERY STOCK, SHADE TREES AND ORNAMENTALS

Agway Kelthane 1.5EC controls two-spotted mite (red spider), European red mite, Spruce mite and other mites which are common in greenhouse and outdoor plants (roses, flowers, ornamentals, shade trees and evergreens).

Winter foliage may be more sensitive than summer foliage. Do not use on sensitive varieties or tender foliage.

### CLOVER MITES IN HOMES AND BUILDINGS

For control of clover mites in homes, spray walls, foundations, windowsills and plants and lawns at base of infested building.

### LAWN AND TURF GRASSES

To control clover, bermuda and brown mites when they appear use two tablespoons per gallon. Apply 2½ gallons of dilute spray per 1000 sq. ft. Repeat application in 10-14 days if necessary.

### VEGETABLES

(PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS)

For control of European red, clover (brown almond), two-spotted (red spider), Schoene, McDaniel, Pacific Willamette, peach silver, plum, nursery and yellow mites, apply at petal fall to first cover spray or when mites appear. Repeat if necessary. Wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes, figs within 7 days of harvest. Do not apply to other fruits within 14 days of harvest.

GRAPES, CHESTNUTS, FILBERTS, HICKORY NUTS, PECANS, WALNUTS, ALMONDS

For control of Brown Almond (clover), two spotted (red spider), European red, Pacific, McDaniel, Willamette and yellow mites apply just before bloom or whenever mites appear. Do not apply on almonds after nut formation. Do not apply on hickory nuts, chestnuts and walnuts within 14 days of harvest. Do not apply on grapes within 7 days of harvest.

### STRAWBERRIES

For control of two-spotted mite (red spider), strawberry mite (Atlantic mite), and six-spotted mite, use 2-4 teaspoons per gallon of water. Drench plant thoroughly at 10-20 day intervals when new growth starts. Use as a spot treatment during season. Do not apply within 2 days of harvest.

## CAUTION:

### KEEP OUT OF REACH OF CHILDREN

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Harmful if swallowed. Avoid inhalation of spray mist. Do not get in eyes, on skin or on clothing. In case of skin contact, wash with soap and water. Wash thoroughly after handling. Avoid contamination of food and feed.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from

#### ORNAMENTALS

Agway Kelthane 1.5EC controls two-spotted mite (red spider), European red mite, Spruce mite and other mites which are common in greenhouse and outdoor plants (roses, flowers, ornamentals, shade trees and evergreens).

Winter foliage may be more sensitive than summer foliage. Do not use on sensitive varieties or tender foliage.

#### CLOVER MITES IN HOMES AND BUILDINGS

For control of clover mites in homes, spray walls, foundations, windowsills and plants and lawns at base of infested building.

#### LAWN AND TURF GRASSES

To control clover, bermuda and brown mites when they appear use two tablespoons per gallon. Apply 2½ gallons of dilute spray per 1000 sq. ft. Repeat application in 10-14 days if necessary.

#### VEGETABLES

For control of two-spotted mite (red spider), Pacific, Tropical and Tomato russet mites on beans, cucumbers, cantaloupes, melons, watermelons, pumpkins, squash (winter and summer), tomatoes, eggplants and peppers. Do not apply on beans within 7 days of harvest. Do not feed treated fresh hay or spent hay to dairy or meat animals. Do not apply on other vegetables within 2 days of harvest.

#### FRUITS (APPLES, PEARS, CRABAPPLES, QUINCES)

For control of European red mite, clover mite, two-spotted (red spider), Schoene, McDaniel, yellow (carpini), Willamette, Pacific and apple rust mites, apply in second cover spray or whenever mites appear. Do not apply within 7 days of harvest.

#### STRAWBERRIES

For control of two-spotted mite (red spider), strawberry mite (Atlantic mite), and six-spotted mite, use 2-4 teaspoons per gallon of water. Drench plant thoroughly at 10-20 day intervals when new growth starts. Use as a spot treatment during season. Do not apply within 2 days of harvest.

### CAUTION:

### KEEP OUT OF REACH OF CHILDREN

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Harmful if swallowed. Avoid inhalation of spray mist. Do not get in eyes, on skin or on clothing. In case of skin contact, wash with soap and water. Wash thoroughly after handling. Avoid contamination of food and feed.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with wastes or by burning. Stay away from smoke.

**DISCLAIMER:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-449

AGWAY, INC.

FERTILIZER-CHEMICAL DIVISION, BOX 1333  
SYRACUSE, NEW YORK 13201

**ACCEPTED**

7-16-76

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR EQUIPMENT WHICH IS REGISTERED  
UNDER NO. 8590-449