

ADVERTISING AND SALES PROMOTION SERVICES

1284
PM 12
8590 149
COPY

FOR:

KELTHANE (R) 1.5EC

DATE: 9/19/84

Net Contents

pt. qt.

AGWAY

KELTHANE (R) 1.5EC

Agricultural miticide controls mites on vegetables, fruits, shade trees, ornamentals, flowers, and controls clover mites in homes.

ACTIVE INGREDIENT :

1, 1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	18.5%
Xylene range aromatic solvent	73.0%

INERT INGREDIENTS:

8.5%

TOTAL 100.0%

(R) Kelthane, registered trademark of Rohm & Haas Co.

KEEP OUT OF REACH OF CHILDREN (12 pt)

CAUTION (18 pt)

See additional precautionary statements on back panel.

AGWAY INC.
Chemical Division
P. O. Box 4933
Syracuse, New York 13221

EPA Reg. No. 8590-449

EPA Est. No. 8590-PA-1

OK BY

EC
NOV 05 1984
8590-449

ADVERTISING AND SALES PROMOTION SERVICES

10/19/84

FOR:

KELTHANE R 1.5EC

DATE: 9/19/84

((BACK PANEL))

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

STORAGE & DISPOSAL (12 pt)

Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals.

Do not reuse empty container. Wrap container and put in trash.

Use two teaspoons of Kelthane 1.5EC in one gallon of water unless otherwise noted. Spray both upper and lower plant surfaces when mites first appear. Repeat when necessary.

FLOWERS, NURSERY STOCK, SHADE TREES AND ORNAMENTALS

Agway Kelthane 1.5EC controls two-spotted mite (red spider), European red mite, Spruce mite and other mites which are common in greenhouse and outdoor plants (roses, flowers, ornamentals, shade trees and evergreens).

Winter foliage may be more sensitive than summer foliage. Do not use on sensitive varieties or tender foliage.

CLOVER MITES IN HOMES AND BUILDINGS

For control of clover mites in homes, spray walls, foundations, windowsills and plants and lawns at base of infested building.

LAWN AND TURF GRASSES

To control clover, bermuda and brown mites when they appear use two tablespoons per gallon. Apply 2 1/2 gallons of dilute spray per 1000 sq. ft. Repeat application in 10-14 days if necessary.

VEGETABLES

For control of two-spotted mite (red spider), Pacific, Tropical and Tomato russet mites on beans, cucumbers, cantaloupes, melons, watermelons, pumpkins, squash (winter and summer), tomatoes, eggplants and peppers. Do not apply to cucumbers and tomatoes within 21 days of harvest. Do not apply on beans within 7 days of harvest. Do not

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

308 u
COPY

FOR:

KELTHANE R 1.5EC

DATE: 9/19/84

feed treated fresh hay or spent hay to dairy or meat animals. Do not apply on other vegetables within 2 days of harvest.

FRUITS (APPLES, PEARS, CRABAPPLES, QUINCES)

For control of European red mite, clover mite, two-spotted (red spider), Schoene, McDaniel, yellow (carpini), Willamette, Pacific and apple rust mites, apply in second cover spray or whenever mites appear. Do not apply within 7 days of harvest.

(PEACHES, APRICOTS, NECTARINES, CHERRIES, PLUMS, PRUNES, FIGS)

For control of European red, clover (brown almond), two-spotted (red spider), Schoene, McDaniel, Pacific Willamette, peach silver, plum, nursery and yellow mites, apply at petal fall to first cover spray or when mites appear. Repeat if necessary. Wait 30 days to avoid excessive residues. Do not apply to cherries, plums, prunes, figs within 7 days of harvest. Do not apply to other fruits within 14 days of harvest.

GRAPES, CHESTNUTS, FILBERTS, HICKORY NUTS, PECANS, WALNUTS

For control of Brown Almond (clover), two-spotted (red spider), European red, Pacific, McDaniel, Willamette and yellow mites apply just before bloom or whenever mites appear. Do not apply on almonds after nut formation. Do not apply on hickory nuts, chestnuts and walnuts within 14 days of harvest. Do not apply on grapes within 7 days of harvest.

STRAWBERRIES

For control of two-spotted mite (red spider), strawberry mite (Atlantic mite), and six-spotted mite, use 2-4 teaspoons per gallon of water. Drench plant thoroughly at 10-20 day intervals when new growth starts. Use as a spot treatment during season. Do not apply within 2 days of harvest.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid inhalation of spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling.

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

410
COPY

FOR:

KELTHANE R 1.5EC

DATE: 9/19/84

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. DO NOT INDUCE VOMITING UNLESS NO OTHER TREATMENT IS AVAILABLE. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin: Remove contaminated clothing and wash affected areas with soap and water. See a physician if irritation persists.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated.

DISCLAIMER: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

OK BY