

PM-1 R-1 8570-455

Agway Livestock & Farm Spray

Permethrin Insecticide

ACCEPTED
JAN 4 1980
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8570-455

stuffed furniture. Do not overspray if products are made of rubber or are rubber coated, to prevent expansion or swelling. Repeat treatments monthly unless the articles are stored in moth-tight containers.

To control infestations of granary weevil, rice weevil, sawtoothed grain beetle, confused flour beetle, rusty grain beetle, red flour beetle, cadellies, meal worms, grain mites, Angoumois grain moth, Indian meal moth and Mediterranean flour moth in food storage areas: If possible, clean out and destroy any infested food products; otherwise, fumigation or other effective treatment should be made using an approved product intended for this purpose. Apply this spray thoroughly to all surfaces of shelves, cabinets, bins, pantries and similar food storage areas with particular attention to cracks, crevices, niches, and other hiding places.

Acceptable for use on dairy and beef cattle, sheep, swine, poultry and horses, in dairy barns, milk rooms, dairies, homes, institutions, restaurants and food processing plants.

Kills and repels exposed or accessible stages of: Roaches, Silverfish, Crickets, Spiders, Ants, Fleas, Cadellies, Flies, House flies, Stable flies, Horse flies, ~~House flies~~, Gnats, Wasps, Hornets, Sand Flies, Mosquitoes, Bedbugs, Meal worms, Meal moths (Indian Meal Moth), Rice Weevil, Spider Beetles, Grain Mites, Cheese mites, Carpet beetles, Angoumois grain moth, Confused flour beetle, Saw Toothed grain beetle, Mediterranean flour moth, Brown Dog ticks in buildings, Granary weevil, Red flour beetle, Clothes moths.

Harmful if swallowed. Avoid contamination of feed and food stuffs.

The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Feeds should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Remove pets, birds, and cover fish aquaria before spraying.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

FOG OR VAPOR SPRAY:

To kill accessible or exposed stages of insects named on this label (excepting clothes moths, bedbugs, roaches, spiders, crickets and carpet beetles). Apply using conventional mechanical thermal or compressed air equipment. Fill the entire area with mist using 3 to 6 ounces per 1000 cubic feet of space. Repeat as needed.

OUTDOOR ADULT MOSQUITO CONTROL:

Apply as a fine mist to thoroughly treat the entire area. For best results treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Shrubby and vegetation around low marshy areas, stagnant pools or ponds and shore lines may be treated. Application to any body of water is prohibited. Apply at a rate of 1/2 gallon per acre (1 1/2 fluid ounces per 1000 square feet).

Agway Livestock & Farm Spray can be used in thermal, mechanical or compressed air spray equipment. Calibration of conventional truck mounted mist blowing equipment should be based on traveling 5 mph to treat a 300 foot swath to deliver 1 1/2 gallons per minute.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given here-with.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.



Agway Inc.
 Chemical Division, Box 4000

BEST AVAILABLE COPY

A ready-to-use Pyrenone Insecticide

BEST

ACTIVE INGREDIENTS:

Pyrethrins	0.30%
Piperonyl Butoxide, Technical*	2.50%
Petroleum Distillate	97.20%
	<u>100.00%</u>

*Equivalent to 2.0% of (butylcarbityl 6-propylpiperonyl) ether and 0.5% of related compounds.

DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide spray mixture or rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill or bury in a safe place.

Consult federal, state or local disposal authorities for approved alternative procedures.

DAIRY & BEEF CATTLE, SHEEP, SWINE, POULTRY, AND HORSES. To protect animals from horn flies, house flies, Mosquitoes, and gnats, apply a light mist sufficient to wet the tips of hair. To control stable flies, horse flies, deer flies, apply 2 ounces per adult animal, sufficient to wet the hair but not to soak the hide. Repeat treatment once or twice daily or at intervals necessary to give continued protection.

To control face flies, moisten cloth with Agway Livestock and Farm Spray and wipe animal facial area.

To control blood-sucking lice, apply the spray lightly to infested areas, using a stiff brush to get the spray to the base of the hair. Use 2 ounces or less per animal. Repeat every two or three weeks as required.

SPACE SPRAY

For rapid control of flies, fruit flies, mosquitoes, gnats, small flying moths, sand flies, wasps and hornets: Close doors and windows and shut off ventilating systems. Apply using a good machine adjusted to deliver a fine mist. Direct spray toward the upper portions of the room to fill the entire area with mist, using 1 to 2 ounces per 1000 cubic feet of space. Vacate the treated area and ventilate before reoccupying. Keep the area closed for ten minutes after treating. Repeat application when reinfestation occurs.

SURFACE APPLICATION

To control roaches, silverfish, crickets, and spiders: Apply as a course wetting spray directed into locations where these insects may hide, behind baseboards, moldings, cabinets, shelves, etc. door and window frames, and clutter.

stuffed furniture. Do not overspray if products are coated, to prevent expansion or swelling. Repeat articles are stored in moth-tight containers.

To control infestations of granary weevil, rice confused flour beetle, rusty grain beetle, red flour grain mites, Angoumois grain moth, Indian meal moth in food storage areas: If possible, clean out products; otherwise, fumigation or other effective an approved product intended for this purpose. Pay particular attention to cracks, crevices, niches, and

Acceptable for use on dairy and beef cattle, sheep, swine, poultry and horses, in dairy barns, milk rooms, dairies, homes, institutions, restaurants and food processing plants.

Kills and repels exposed or accessible stages of: Roaches, Silverfish, Crickets, Spiders, Ants, Fleas, Caddises, Flies, House flies, Stable flies, Horse flies, Horn flies, Lice, Gnats, Wasps, Hornets, Sand Flies, Mosquitoes, Bedbugs, Meal worms, Meal moths (Indian Meal Moth), Rice Weevil, Spider Beetles, Grain Mites, Cheese mites, Carpet beetles, Angoumois grain moth, Confused flour beetle, Saw Toothed grain beetle, Mediterranean flour moth, Brown Dog ticks in buildings, Granary weevil, Red flour beetle, Clothes moths.

Harmful if swallowed. Avoid contamination of feed and food stuffs.

The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Feeds should be removed or covered before treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Remove pets, birds, and cover fish aquaria before spraying.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish. Keep out of water.

FOG C
To kill
insects
clothes
ders.
Apply
therma
Fill the
6 ounce
Repea

OUTDO
CONT
Apply
the en
when
5 mile
vegeta
Shrub
marsh
and st
cation
ited A
(1 1/2 fl

Agway
used
pressu
tion o
mist
based
foot s
minut

DISC
of use
accor
with

