



# DICHLONE - SULFUR 1.5 - 30D

RECOMMENDED FOR THE CONTROL OF FUNGUS DISEASES ON APPLES, CHERRIES, PLUMS AND PEACHES. THIS DUST IS COMPATIBLE WITH THE DRY POWDERS OF PARATHION, THIRAM, FERBAM, CAPTAN, AND ZINEB.

ACTIVE INGREDIENTS:	
Dichlone (2,3-dichloro - 1,4-naphthoquinone) . . . . .	1.5%
Sulfur . . . . .	30.0%
INERT INGREDIENTS: . . . . .	68.5%
	100.0%

## DIRECTIONS

READ THESE ENTIRE DIRECTIONS AND WARRANTY OF SALE. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

**APPLES** - To control Apple Scab and Powdery Mildew apply this dust at the rate of 60 to 80 pounds per acre. Begin applications at first sign of green tissue and repeat applications every 5 to 7 days up through the first cover spray (7 days after petal fall). Do not use this dust after first cover. McIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulfur applications under certain conditions.

**PEACHES** - To control Peach Scab, Brown rot of fruit, Brown rot blossoms blight and Cercospora leaf spot, apply dust at the rate of 60 to 80 pounds per acre. Applications should begin at the pre-bloom stage and continue every 5-7 days through the bloom, petal fall and first four cover applications. Dust may be used every 3-4 days if rainy weather prevails. For Brown Rot and Leaf Spot this dust may be used at weekly intervals to within 7 days of harvest.

**CHERRIES** - For the control of Brown rot of fruit, Brown rot blossom blight and cercospora leaf spot apply as directed on peaches. Do not apply within 3 days of harvest.

**PLUMS** - For the control of Brown rot of fruit, Brown rot blossom blight and phytoxic leaf spot, apply as indicated on peaches. Do not apply within 3 days of harvest.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid inhalation or contact with skin. May cause skin irritation. In case of contact with skin, wash thoroughly with soap and warm water. For contact with eyes, flush with plenty of water and get medical attention.

Wash hands and face with soap and warm water after using and before eating or smoking. Avoid contamination of feed and food stuffs.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Sulfur may cause severe fruit and foliage injury to certain crops. Consult with your State Agricultural Experiment Station or your State Agricultural Extension Service for additional information consistent with this label's cautions and directions. Sulfur may burn foliage when temperature is high. Do not make sulfur applications at such times. Do not use within 2 weeks of an oil spray treatment. When growing crops for processing, consult the processor before applying sulfur.

**WARRANTY:** Seller warrants that the product conforms to its chemical description and is reasonable fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

**NOTE:** This pesticide is to be sold ONLY in this original, unbroken package.

EPA Reg. No. 8590-461

EPA Est. No. 8590-PA-1

AGWAY INC.,  
FERTILIZER-CHEMICAL DIVISION  
BOX 1333, SYRACUSE, NEW YORK 13201

NET WEIGHT FIFTY POUNDS