



BLUE RIBBON RAT PACK

20016 16F1
EPA REG
1 03 01

A TOX-HID* COATED WARFARIN PELLETTED BAIT.

Kills Norway Rats, Roof Rats and House Mice.

For Indoor Use Only

Each particle of the Warfarin in this product has been specially coated with a material that reduces detection of the Warfarin by the rodents at the time the bait is consumed. The coating may help assure greater acceptance of the bait in cases where rodent awareness of uncoated Warfarin is responsible for low bait acceptability.

ACTIVE INGREDIENT:

Warfarin (3-alpha-Acetylbenzyl)-4-hydroxycoumarin

INERT INGREDIENTS: 99.975%

TOTAL 100.000%

*TOX-HID — Wisconsin Alumni Research Foundation — Patent Pending. TOX-HID — Trademark of Wisconsin Alumni Research Foundation.

CAUTION
KEEP OUT OF REACH OF CHILDREN.

PRECAUTIONARY STATEMENTS:

Hazard to Humans & domestic animals:

CAUTION: Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K₁, combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdose of bishydroxycoumarin.

ENVIRONMENTAL HAZARD: Keep out of Lakes, Streams or Ponds.

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS — For control of Norway rats, roof rats and House mice in homes, industrial and agricultural buildings, and similar man-made structures. For indoor use only. Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals, or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS — Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Place 1 to 4 packs per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Open pack and apply ¼ to ½ ounce (½ level tablespoons) of bait at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

Do not reuse empty container.

Wrap container and put in trash collection.

EPA Reg No 8590-475

EPA Est No 8590-NY-1

AGWAY INC. — CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

A 12/80

NET CONTENTS: 4 OZ.