



COMPANY

NUMBER 8590

PRODUCT

SERIAL NO. 487

LABEL

APPROVAL

DATE

6-5-96

JUN 05 1996

Beverly Randall
Agway Inc.
c/o Universal Cooperatives, Inc.
P.O. Box 460
Minneapolis, Minnesota 55440

Dear Ms. Randall:

Subject: Deleted uses
Agway 25% Methoxychlor Spray
EPA Registration No. 8590-487
Amendment application of October 11, 1995

The amendment referred to above, submitted in connection with registration under FIFRA, is acceptable provided that you:

The direction for use to control mosquitoes has been deleted from the label. This use is no longer being supported, therefore you must delete the following statement wherever it appears on the labeling:

"Can also be use to control mosquitoes outdoors."

Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide Rodenticide Branch
Registration Division (7505C)

Enclosure

((FRONT PANEL))

NET CONTENTS
1 pt.

AGWAY
25% METHOXYCHLOR SPRAY

An insecticide emulsifiable concentrate containing 2 lbs Methoxychlor per gallon for home and garden use only.

For control of various insects on fruits, vegetables, ornamentals, and shade trees. Can also be used to control mosquitoes outdoors.

ACTIVE INGREDIENTS:

Methoxychlor*:	1,1,1-Trichloro-2,2-bis (p-methoxyphenyl) ethane	25.0%
----------------	---	-------

INERT INGREDIENTS**:		75.0%
	TOTAL	<u>100.0%</u>

*Equivalent to 22% 1,1,1-trichloro-2,2-bis (p-methoxyphenyl) ethane, 3% other isomers and related compounds.

**Contains xylene range aromatic solvent.

KEEP OUT OF REACH OF CHILDREN. (12 pt)

CAUTION (18 pt)

See additional precautionary statements on side panel.


AGWAY INC.
P.O. BOX 4741
SYRACUSE, NY 13221

EPA Reg. No. 8590-487
EPA Est. No. 8590-PA-1

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUN 05 1996

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
8590-487



((REAR PANEL))

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. Do not induce vomiting unless no other treatment is available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin: Remove contaminated clothing and wash affect areas with soap and water. See a physician if irritation persists.

If in eyes: Flush eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS:

This product is toxic to fish and aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water or wetlands (swamps, bogs, marshes & pot-holes). Do not contaminate water when disposing of equipment washwaters.

Do not use or store near heat or open flame. Do not apply within 14 days of sulfur or sulfur containing products.

STORAGE DISPOSAL (12 pt)

Store unused product in original container only. Store in a cool, dry area, preferably locked and out of reach of children.

Do not reuse empty container. Wrap container and put in trash collection.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE.

These directions are for home garden use only, not for large scale agricultural applications. Thoroughly mix 1 1/2 to 2 tablespoons of this product in 1 gallon of water, unless otherwise directed. Spray thoroughly to cover all plant surfaces. Begin applications at first sign of infestations; on fruits, begin at petal fall. Repeat at 7 to

14 days intervals, as needed, to maintain effective control. Do not spray food crops within the number of days of harvest shown in parentheses for each crop.

For control of various insects on fruits, vegetables, ornamentals, and shade trees. Can also be used to control mosquitoes outdoors.

Food utensils (teaspoons, tablespoons) should not be used for food purposes after use in measuring pesticides.

FRUIT TREES

On apples (7), apricots (21), cherries (7), nectarines (21), peaches (21), pears (7), plums (7), prunes (7), quinces (7). Controls apple maggot, cankerworm, cherry fruit fly, cherry fruitworm, codling moth, Japanese beetle, leafhopper, pear slug, plum curculio, tent caterpillar.

VEGETABLES

On asparagus (3), beans (3), beets* (7), broccoli (14), brussels sprouts (14), cabbage (3), cantaloupe (7), carrots* (7), cauliflower (7), collards (14), corn** (7), cucumber (1), eggplant (1), kale (14), kohlrabi (1), lettuce (4), melons (7), peas** (7), peppers (1), potatoes (0), pumpkins (1), radishes (7), rutabagas (7), spinach (14), squash (1), summer squash (1), tomatoes (7), turnips* (7). Controls asparagus beetle, bean leaf beetle, blister beetle, colorado potato beetle, corn earworm, cross-striped cabbageworm, cucumber beetle, fall armyworm, flea beetle, imported cabbage worm, Japanese beetle, leafhopper, melon worm, Mexican bean beetle, pea weevil, rose chafer, squash vine borer, and tomato hornworm.


*Beets, carrots, turnips: If tops are used for food, do not spray within 14 days of harvest.

**Corn: Do not feed treated corn to livestock.

***Peas: Do not graze or feed treated pea vines or pea hay to livestock.

SMALL FRUITS AND BERRIES

On blackberries (14), blueberries (14), currants (14), gooseberries (14), grapes (14), loganberries (14), raspberries (14), and strawberries (14). Controls cherry fruitworm, grape berry moth, grape leaf skeletonizer, Japanese beetle, leafhopper, rose chafer, San Jose scale (crawlers), spittlebug, and strawberry weevil.



TO CONTROL INSECTS ON SHADE TREES, FLOWERS, OR ORNAMENTALS: Dilute 3/4 - 1 1/2 teaspoons per gallon of water to control blister beetle, cankerworm, cucumber beetle, Eastern tent caterpillar, elm leaf beetle, fall webworm, fall armyworm, flea beetle, forest tent caterpillar, gypsy moth, inchworm, Japanese beetle, leafhopper, lace bug, leaf roller, May beetle, rose chafer, rose slug, sawfly, spittlebug, tent caterpillar, and tussock moth.

DISCLAIMER: Buyer assumes all risks of use, storage, or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold only in this original, unbroken package.

