

AGWAY DIAZINON® 50W

For control of certain insects on fruit, nuts, vegetables, field crops and ornamentals

ACTIVE INGREDIENT

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 50.00%

INERT INGREDIENTS 50.00%

TOTAL 100.00%

Diazinon® Trademark of Ciba-Geigy

WARNING: KEEP OUT OF REACH OF CHILDREN

See additional precautionary statements

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. See separate sheet for directions.

STORAGE & DISPOSAL

Do not reuse empty container

Not for use or storage in the home

Do not contaminate water, food or feed by storage, cleaning of equipment or disposal. Open dumping is prohibited.

Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets. Do not permit children or pets to go on to sprayed grass until spray has completely dried.

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled — Remove victim to fresh air. Apply respiration if indicated.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes for at least 15 minutes with soap and water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries (except for effluence from treated watercross beds). Do not apply to water that will be used for recreational purposes and human and livestock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. This product must not be used in areas where impact on threatened endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment areas. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

DISCLAIMER: Buyer assumes all risks of use, storage, or handling not in strict accordance with directions given hereon.

NOTE: This pesticide is to be stored only in this original unbroken package.

AGWAY INC., HEMET, CALIF. 92343

BOX 493, SYRACUSE, N.Y. 13221

EPA Reg. No. 8590-494

EPA Reg. No. 8590-PA-1

NET CONTENTS 5 LBS.

81

SEW

SEW



Phly-N-Worm 4.4

Aids in preventing the breeding of horn flies and face flies in the manure of treated cattle, when used as directed.

Aids in the control of stomachworms (*Haemonchus*, *Ostertagia*, and *Trichostrongylus* spp.); nodular worms (*Oesophagostomum* spp.); large-mouth bowel-worms (*Chabertia* spp.); and hookworms (*Bunostomum* spp.) in cattle when fed in accordance with the directions for use.

ACTIVE INGREDIENT

Phenothiazine.....4.4%

INERT*INGREDIENTS.....95.6%

TOTAL

100.00%

*Refers only to ingredients which are not pesticides.

Each lb. DALE ALLEY PHLY-N-WORM 4.4 contains 20 g. Phenothiazine.

FEED INGREDIENTS

Processed Grain By-Products, Calcium Carbonate and Mineral Oil

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

32695-7

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

HAZARDS TO DOMESTIC ANIMALS

Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. Do not administer to animals showing symptoms of acute respiratory conditions.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

See Reverse Side For Directions For Use and Storage and Disposal



NET WEIGHT 10 LBS.

Manufactured by

DALE ALLEY CO.

St. Joseph, Missouri 64502

EPA Reg. No. 32695-7

EPA Est. No. 10543-MO-01

19 LOT NO.

DIRECTIONS FOR USE

It is in violation of federal law to use this product in a manner inconsistent with its labeling.

Mix 1 pound of **ALLEY Phly-N-Worm 4.4** in the daily feed, minerals or salt** of 10 adult cattle to provide the 2 grams Phenothiazine intended.

**Self-feed continuously as the sole source of salt.

Younger cattle: Follow the feeding chart below to provide $\frac{1}{4}$ gram Phenothiazine per 100 pounds body weight per head daily.

AVERAGE BODY WEIGHT OF YOUNGER CATTLE	Lbs. ALLEY Phly-N-Worm 4.4		
	per 100	1000	2000 feedings
200 lbs.	2 $\frac{1}{4}$	25	50
400 lbs.	5	50	100
500 lbs.	6 $\frac{1}{4}$	62 $\frac{1}{2}$	125
600 lbs.	7 $\frac{1}{2}$	75	150
700 lbs.	8 $\frac{1}{4}$	87 $\frac{1}{2}$	175

Continuous year round low-level phenothiazine treatment, in conjunction with good herd management practices, will control the specified parasites and will help prevent reinfection.

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

Cattle should be given access to feed, minerals or salt containing **ALLEY Phly-N-Worm 4.4** for several weeks prior to the appearance of adult horn flies in the spring, and should be fed a **ALLEY Phly-N-Worm 4.4** supplement continuously until flies disappear in the fall. This helps prevent the breeding of horn flies and face flies but has no effect upon existing adult fly populations. Since horn flies over-winter in the pupal state, all cattle should be sprayed when the flies first appear. Spraying kills the flies emerging from last year's pupae and will protect cattle for about two weeks until the phenothiazine can take effect.

Since face flies migrate from farm to farm it is essential that the larvicide program be carried out on an area-wide basis to assist in the reduction of the adult face fly population.

STORAGE AND DISPOSAL

Prohibitions:

Do not contaminate water or food by storage and disposal.

Pesticide Disposal:

Feed supplements containing Phly-N-Worm 4.4 which cannot be used should be buried in a safe place away from water supplies.

Container Disposal:

Do not reuse empty container. Wrap container and put in trash collection.