

ACCEPTED

AUG 27 1982

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended for the pesticide
registered under
EPA Reg. No. 8590-495

AGWAY THIRAM 75W

FRUIT, VEGETABLE AND TURF FUNGICIDE

For the control of Dollar Spot, Brown Patch, and Snow Mold.

ACTIVE INGREDIENT	
Thiram (Tetramethylthiuram Disulfide)	75%
INERT INGREDIENTS	25%
TOTAL	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bag in a sanitary landfill or by incineration if allowed by State and local authorities

DIRECTIONS FOR USE AS A SPRAY ON FRUITS & VEGETABLES.

ALL RECOMMENDATIONS BASED ON 100 GALLONS OF SPRAY EXCEPT AS NOTED.

APPLES: Scab — 1.3 to 1.7 pounds in pre-bloom, calyx, early cover sprays; and later cover sprays

Rest — Use scab recommendations through first cover and include a special bloom spray of 0.65 pound.

Other Diseases — Use scab recommendations as long as any one or more of the following diseases may appear — Apple Blotch, Bitter Rot, Black Pox, Black Rot (fruit and foliage), Botryosphaeria Spot, Brooks Spot, Fly Spack and Spotty Blotch.

CELERY: Early Blight (Cercospora), **Late Blight** (Septoria) and **Damping-Off Plant Beds** — 1.3 pounds. Apply 15 to 25 gallons/1200 sq. ft. at 3-day intervals (Plant beds, only)

PEACHES: Brown Rot — For blossom blight, apply 1.3 to 1.7 pounds at 3 to 4 day intervals during bloom, timed to keep newly opened blossoms covered. For fruit rot, apply at petal-fall and two weeks later. Make one or more additional applications at 7-day intervals beginning about one month before harvest, or earlier if rot is apparent on green fruit

Rhizopus Rot — Apply as directed for fruit phase of brown rot

Scab — Apply at petal-fall and repeat 3 to 6 times at 10 to 14-day intervals or until disease is well under control

Do not apply later than 7 days before harvest

STRAWBERRIES: Gray Mold (Botrytis) — As a spray, use 1.7 to 2.2 lbs. per 100 gallons of water applying at rates up to 200 gallons per acre. As a dust use a mixture containing 6.5% to 8.7% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite applying at the rate of 40 lbs. per acre. Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results

If applied within 3 days of picking, remove residues by washing

TOMATOES: Botrytis Gray Mold and Rhizoctonia Fruit Rot — 1.3 pounds when plants are set in the field and repeat at 3 to 5-day intervals. If late blight, gray leaf spot or other foliage diseases are present, use combination of 0.9 pounds Thiram 75W with 1 1/2 lbs. of either Maneb 80% or Zineb 75% wettable powder fungicides. Make first application when fruits first are visible on the first cluster (late varieties) or just after blossoming (early varieties). Continue at 7 to 10 day intervals, depending upon weather, until 6 sprays have been applied. When tank mix is used, do not apply combination within 5 days of harvest.

DIRECTIONS FOR USE AS A SEED AND SOIL TREATMENT FOR CONTROL OF DAMPING-OFF, SEED DECAY, SEEDLING BLIGHTS, ONION SMUT AND SWEET POTATO STEM ROT

Apply at recommended dosages using the dry powder as received or mixed with water to form a slurry. Treated seed should be labeled, "Do not use treated seed for food, feed or oil"

BEANS, Lima — 2.2 oz./100 pounds applied dry

BEANS, String — 1.3 oz./100 pounds applied as a slurry

BEANS, Snap — 1.3 oz./100 pounds applied dry or as a slurry

ONIONS — As a dust furrow treatment — 1.3 pounds/acre mineral soils (14,500 linear feet of row) or 2 1/4#/acre on muck soils (14,500 linear feet of row) (dry bulb)

SOYBEANS — 1.3 oz./100 pounds applied dry or as slurry

SWEET POTATOES — 1.0 pounds/7 1/2 gal. water applied as a preplanting root dip for 1/2 minute. Plant sprouts promptly after treatment

At planting time only. Do not use treated roots for food or feed

Agway Thiram 75W prevents infestation of common turf diseases such as Dollar Spot, and Brown Patch if sprayed at regular weekly intervals throughout the summer season

Agway Thiram 75W will control Dollar Spot and Brown Patch when applied at a higher dosage rate to turf already infested with these diseases

Brown Patch, Dollar Spot — Preventive Treatment. Apply in early spring before disease symptoms appear, then follow a regular 7 to 10 day spray schedule throughout the season. Use 2 1/2 to 3 oz. and apply in sufficient water to cover 1,000 sq. ft.

Curative (eradicator) treatment. At first sign of disease apply 4 1/2 to 5 oz. in water per 1,000 sq. ft. Repeat at 3 to 5 day intervals. When disease is controlled, follow the preventive spray schedule

Snow Mold — Use 8 oz. per 1,000 sq. ft. applied immediately before expected permanent snow cover. If possible, make repeat applications following each thaw using 3 oz. per 1,000 sq. ft. Follow with a spring application of 3 oz. per 1,000 sq. ft. as soon as snow melts

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION May irritate eyes, nose, throat, & skin. Harmful if inhaled or swallowed. Do not get in eyes, on skin or clothing. Wash thoroughly after handling

STATEMENT OF PRACTICAL TREATMENT

In case of contact wash exposed areas of contact with water to prevent irritation. For eyes, get medical attention

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes

DISCLAIMER. Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith

NOTE. This pesticide is to be sold ONLY in this original unbroken package

EPA Reg. No. 8590-495

EPA Est. No. 8590-PA-1



AGWAY INC., CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

NET CONTENTS 3 LBS.

A 7/82