

Reg # 8590-495

1652
12001

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY THIRAM 75W

DATE: 4/1 /85

AGWAY THIRAM 75W Net Wt. 3#
FRUIT, VEGETABLE AND TURF FUNGICIDE

For the control of Dollar Spot, Brown Patch, and Snow Mold.

Active Ingredient:

Thiram (Tetramethylthiuram Disulfide)	75%
INERT INGREDIENTS:	25%
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)


See additional precautionary statements on side panel

EPA Reg. No. 8590-495

EPA Est. No. 8590-PA-1

Agway Inc.
Chemical Division
Box 4933
Syracuse, New York

13221



OK BY

SHEET NO.

1-8

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY THIRAM 75W

DATE: 4/16/85

((((REAR PANEL)))

FILE NO. CHARACTER
DATE NO. CHARACTER

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE AS A SPRAY ON FRUITS & VEGETABLES.

ALL RECOMMENDATIONS BASED ON 100 GALLONS OF SPRAY EXCEPT AS NOTED.

APPLES: Scab - 1.3 to 1.7 pounds in pre-bloom, calyx, early cover sprays; and later cover sprays
Rust - Use scab recommendations through first cover and include a special bloom spray of 0.65 pound.

Other Diseases - Use scab recommendations as long as any one or more of the following diseases may appear - Apple Blotch, Bitter Rot, Black Pox, Black Rot (fruit and foliage). Botryosphaeria Spot, Brooks Spot, Fly Speck and Sooty Blotch.

CELERY: Early Blight (Cercospora), Late Blight (Septoria) and Damping-Off Plant Beds - 1.3 pounds
Apply 15 to 25 gallons/1200 sq. ft. at 3-day intervals. (Plant beds, only)

PEACHES: Brown Rot - For blossom blight, apply 1.3 to 1.7 pounds at 3 to 4 day intervals during bloom, timed to keep newly opened blossoms covered. For fruit rot, apply at petal-fall and two weeks later. Make one or more additional applications at 7-day intervals beginning about one month before harvest, or earlier if rot is apparent on green fruit.

Rhizopus Rot - Apply as directed for fruit phase of brown rot.

Scab - Apply at petal-fall and repeat 3 to 6 times at 10 to 14-day intervals or until disease is well under control. Do not apply later than

Q'RAY

SHEET NO

ERTISING AND SALES PROMOTION SERV!

COPY

FOR:

AGWAY THIRAM 75W

DATE: 4/16/85

7 days before harvest.

PILA - 50 CHARACTER
ELITE - 60 CHARACTER

STRAWBERRIES: Gray Mold (Botrytis) - As a spray, use 1.7 to 2.2 lbs. per 100 gallons of water applying at rates up to 200 gallons per acre. As a dust use a mixture containing 6.5% to 8.7% Thiram (4.88% to 6.5% active) mixed with a suitable diluent such as talc or pyrophyllite applying at the rate of 40 lbs. per acre. Make 3 to 5 applications at 10-day intervals beginning at early blossom stage. Good coverage of strawberry buds, blossoms and developing fruit is required for best results.

If applied within 3 days of picking, remove residues by washing.

TOMATOES: Botrytis Gray Mold and Rhizoctonia Fruit Rot - 1.3 pounds when plants are set in the field and repeat at 3 to 5-day intervals. If late blight, gray leaf spot or other foliage diseases are present, use combination of 0.9 pounds Thiram 75W with 1 1/2 lbs. of either Maneb 80% or Zineb 75% wettable powder fungicides. Make first application when fruits first are visible on the first cluster (late varieties) or just after blossoming (early varieties). Continue at 7 to 10 day intervals, depending upon weather, until 6 sprays have been applied. When tank mix is used, do not apply combination within 5 days of harvest.

Do not rotate treated crop with crops other than those with registered Thiram uses.

Do not reenter treated areas within 24 hours following application unless protective clothing is worn.

ORBY

SHEET NO.

ERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY THIRAM 75W

DATE: 4/16/85

DIRECTIONS FOR USE AS A SEED AND SOIL TREATMENT
FOR CONTROL OF DAMPING-OFF, SEED DECAY, SEEDLING
BLIGHTS, ONION SM'IT AND SWEET POTATO STEM ROT.

FILE NO. CHARACTER
ELITE - 00 CHARACTER

Apply at recommended dosages using the dry powder
as received or mixed with water to form a slurry.
Treated seed should be labeled, "Do not use
treated seed for food, feed or oil."

BEANS, Lima - 2.2 oz./100 pounds applied dry.

BEANS, String - 1.3 oz./100 pounds applied as
a slurry.

BEANS, Snap - 1.3 oz./100 pounds applied dry or
as a slurry.

ONIONS - As a dust furrow treatment - 1.3 pounds
/acre mineral soils (14,500 linear feet of row)
or 2 3/4#/acre on muck soils (14,500 linear
feet of row). (dry bulb)

SOYBEANS - 1.3 oz./100 pounds applied dry or as
slurry.

SWEET POTATOES - 1.0 pounds/7 1/2 gal. water
applied as a preplanting root dip for 1/2 minute.
Plant sprouts promptly after treatment.

At planting time only. Do not use treated roots
for food or feed.

Agway Thiram 75W prevents infestation of common
turf diseases such as Dollar spot, and Brown
Patch if sprayed at regular weekly intervals
throughout the summer season.

Q'KBY

Agway Thiram 75W will control Dollar Spot and
Brown Patch when applied at a higher dosage
rate to turf already infested with these diseases.

Brown Patch, Dollar Spot - Preventive Treatment.
Apply in early spring before disease symptoms
appear, then follow a regular 7 to 10 day spray
schedule throughout the season. Use 2 1/2 to
3 oz. and apply in sufficient water to cover
1,000 sq. ft.

SHEET NO.

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY THIRAM 75W

DATE: 4/16/85

Curative (eradicator) treatment: At first sign of disease apply 4 1/2 to 5 oz. in water per 1,000 sq. ft. Repeat at 3 to 5 day intervals. When disease is controlled, follow the preventive spray schedule.

PICA - 50 CHARACTER
ELITE - 60 CHARACTER

Snow Mold - Use 8 oz. per 1,000 sq. ft. applied immediately before expected permanent snow cover. If possible, make repeat applications following each thaw using 3 oz. per 1,000 sq. ft. Follow with a spring application of 3 oz. per 1,000 sq. ft. as soon as snow melts.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Causes eye & skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap & water after handling. Avoid inhaling spray mist. Do not use or store near food or feed.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, Call a Physician or Poison Control Center immediately. Drink 1 or 2 glasses of water & induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If on skin, remove contaminated clothing & wash affected areas with soap & water. Get medical attention if irritation persists. Wash contaminated clothing before reuse.

G'K BY

If in eyes, flush eyes with plenty of water. Get medical attention if irritation persists.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

ERTISING AND SALES PROMOTION SERVICES

COPY

OR:

AGWAY THIRAM 75W

DATE: 4/16/85

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

During Commercial exposure in seed treatment & loading operations, wear goggles, clean rubber gloves, protective clothing, and a pesticide respirator jointly approved by the Mine Safety & Health Administration (formerly The U.S. Bureau of Mines) and by the National Institute of Occupational Safety & Health under provisions of 30 CFR, Part 11. Wash hands, arms & face thoroughly with soap & water after handling & before eating or smoking. Routinely shower or bathe after work. Wash all clothing with soap & water before reusing.

Consumption of Alcoholic beverages increases the toxic effects of Thiram.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Drift or runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water or wetlands.

STORAGE & DISPOSAL (12 pt.)

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Store unused product in original container only. Store in a cool, dry area preferably locked & out of the reach of children and animals. Not for use or storage in or around the home. Protect from excess heat.

PESTICIDE DISPOSAL: Pesticide Wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your

SHEET NO.

OR:

AGWAY THIRAM 75W

DATE: 4/16/85

State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

RE-ENTRY

Do not enter treated areas without protective clothing until Sprays have Dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

CAUTION

(Insert area or field description) treated with Thiram on (insert date of application). Do not enter treated areas without protective clothing until Sprays have Dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Thiram and describe his condition. For further information see the "Statement of Practical Treatment."

Oral warnings must include the same information that is in the written warnings.

PICA - 50 CHARACTERS
ELITE - 80 CHARACTERS

OK BT

SHEET NO.

7-8

