

Agway Maneb Garden Fungicide

Controls blight, anthracnose, leaf spot and certain other diseases on tomatoes, other vegetables, fruits and flowers.

For Home Garden Use Only.

ACTIVE INGREDIENT	
Maneb (Manganese ethylene bisdithiocarbamate)	80
Manganese equivalent as metallic	16.5
INERT INGREDIENTS	20
	100

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for additional precautionary statements

DIRECTIONS FOR USE: It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THESE ENTIRE DIRECTIONS AND DISCLAIMER. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

STORAGE AND DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

Controls diseases indicated on plants listed

Unless otherwise directed use 1-1 1/2 level tablespoons per gallon of water to make spray mix. Thoroughly spray both upper and under surfaces of foliage. Unless otherwise directed begin when disease first appears. Repeat every 7 to 10 days as needed. Spray more frequently if infestation is severe. When a number in parentheses follows the plant name do not spray that crop within the number of days of harvest shown.

VEGETABLES

Tomatoes (5) — Anthracnose, early and late blight, gray leaf spot (stemphylium), septoria leaf spot. Begin application when tomatoes on first cluster are well-formed. Do not use on young plants in greenhouse.

Beans (4) — Green and dry, Downy mildew, rust. Begin when plants are small. Repeat every 5 to 7 days.

Broccoli (7) — Alternaria leaf spot, downy mildew. If applied within 3 days of harvest remove residues by washing or trimming.

Brussels Sprouts, Cauliflower, Kohlrabi (7) — Alternaria leaf spot, downy mildew.

Cabbage (7) — Alternaria leaf spot, downy mildew. Begin 7 to 10 days after planting. Repeat every 3 to 5 days.

Carrots — Cercospora leaf spot. Start spraying when plants are 6 weeks old. Disease threatens.

Celery (14) — Early and late blight. Start spraying when plants are established in the garden. Remove residues at harvest by stripping, trimming, and washing.

Cucumbers (5), Squash (5) — Anthracnose, downy mildew. Begin when disease first appears or runners start to form.

NET WEIGHT 6 OZ.

Cantaloupe (5), Watermelon (5) — Anthracnose, downy mildew, cercospora leaf spot, downy stem blight. Begin when vines begin to mature. Disease first appears.

Eggplant, Peppers — Anthracnose, blight. Apply as fruit begins to form. Repeat weekly through harvest.

Lettuce (10), Endive (10) — Downy mildew. Remove residues at harvest from head lettuce by stripping and trimming outer leaves, and from leaf lettuce and endive by washing.

Onions — Botrytis leaf blight, downy mildew, purple blotch.

Pumpkins — Angular leaf spot, downy mildew. Use 3 tablespoons per gallon. Spray thoroughly.

Spinach (10) — Downy mildew, white rust. Remove residues of harvest by washing.

Potatoes — Early and late blights. Use 2 to 5 level teaspoons per gallon of water. Apply to 500 sq. ft. as soon as plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals as long as blight threatens. Under severe disease conditions, use the shorter interval or the higher dosage rate.

FLOWERS

Begin spraying when new growth starts or when leaves first unfold. Apply at weekly intervals during season.

Carnations: Leaf spot. **Pansies**: anthracnose. **Dahlias**: botrytis. **Zinnias**: leaf spot. **Flowering Dogwood**: anthracnose. Apply when buds begin to open and again when bracts fall, and 4 weeks later, and again in late summer.

Gladiolus: botrytis, curcularia, Stemphylium leaf spot. **Peony**: botrytis and phytophthora blights. Apply in early spring and early fall to foliage and to soil around plants. Repeat at 7 to 10 day intervals during growing season.

Roses: black spot. **Snapdragon**: rust.

FRUIT

Apples (30) — Black rot, bitter rot, brown rot, bullseye rot, fly speck, rust, scab, sooty blotch, white rot. Begin application in the pre-pink sprays. Repeat in subsequent pre-bloom, bloom and cover sprays in accordance with local state spray schedules.

Do not spray within 30 days of harvest.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS — CAUTION: May cause irritation of nose, throat, eyes and skin. Do not breathe dust or spray mist. May be absorbed through skin.

FIRST AID: In case of skin contact, wash with water. For eyes, wash thoroughly with water and get medical attention.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS: Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may cause chemical changes which will impair the effectiveness of this product and may also generate flammable vapors. Keep container closed when not in use.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOTE: This product is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-500

EPA Est. No. 8590-PA-1

AGWAY — CHEMICAL DIVISION
PO BOX 4000, SYRACUSE, NY 13221

A 7/82