

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

AGWAY® GREENLAWN PLUS®

21-5-7 FERTILIZER WITH DIAZINON INSECTICIDE

Controls White grubs (larvae of Japanese beetle, European Chafer & Southern chafer), sod webworm (lawn moth), chinch bug & other lawn insects as listed.

Feeds the lawn

ACTIVE INGREDIENT

Diazinon: 0,0-Diethyl 0-(2 isopropyl-6-methyl-4

pyrimidinyl phosphorothioate 2.8%

INERT INGREDIENTS\* 97.2%

TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements on back panel.

EPA Reg. No. 8590-502

EPA Est. No. 8590-NY-1

Net wt. 18 Pounds

Treats 5,000 sq. ft.

ACCEPTED  
SEP 27 1994  
Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for two pesticides registered under EPA Reg. No. 8590-502

((BACK PANEL))

Treats 5,000 sq. ft.

1000 sq. ft. = an area 20 ft. x 50 ft.

**Agway Greenlawn Plus**

**21-5-7 Fertilizer With Diazinon Insecticide**

**\*Fertilizer Guaranteed Analysis**

Total Nitrogen (N)	21.00%
6.5 water insoluble nitrogen	
Available phosphoric acid (P2O5)	5.00%
Soluble potash (K2O)	7.00%
Iron (Fe)	0.10%

**Manufactured For:**

**Agway, Inc., Box 4741, Syracuse, NY 13221**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. Avoid repeated contact with skin. If sensitization reactions result, consult a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

**Statement of Practical Treatment**

**If swallowed:** Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

**If inhaled:** Remove victim to fresh air and apply artificial respiration, if indicated.

**If on skin:** Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

**If in eyes:** Rinse eyes with plenty of water and call a physician immediately.

**Note to Physician:** This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

**Environmental Hazards**

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds. Avoid overlapping granules.

For lawns, follow all watering instructions in the directions for use, however, stop watering before puddling occurs. For lawns, do not apply this product to sites occupied by waterfowl, or within 75 ft. of any body of water which may attract waterfowl, such as ponds or lakes.

**Keep out of lakes, streams, ponds, tidal marshes, and estuaries.**

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

---

**DIRECTIONS FOR USE**

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**OVERALL LAWN APPLICATION****Special Precautions To Reduce Risk To Birds**

Birds, especially waterfowl (ducks or geese) feeding or drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

**Storage and Disposal**

**Storage:** Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

**Pesticide Disposal:** Securely wrap original container in several layers of newspaper and discard in trash.

**Container Disposal:** Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

Agway Greenlawn Plus 21-5-7 Fertilizer With Diazinon  
Insecticide rejuvenates starved and insect infested lawn  
by:

- 1) providing a controlled feeding fertilizer which gives a thick green lawn, and
- 2) Controlling the following insect pests which feed on lawns:
  - white grubs (larvae of Japanese beetle, European Chafer, Southern Chafer)
  - sod webworm (lawn moth)
  - armyworms                      chinch bugs              earwigs
  - Bermudagrass mites      clover mites              fleas
  - billbugs                      crickets                      leafhoppers
  - brown dog ticks              Chiggers                      cutworms

How to Use

- 1. Apply uniformly 3.6 lbs./1,000 sq. ft. when insect activity or damage (chewed or brown leaves) are first observed in the lawn. For grub control, treat any time between late July and early October; the next most likely period is Spring.

2. Apply when lawn is dry. The contents of this bag will treat 5,000 sq. ft.
3. Water lawn lightly after application. If treatment is for grubs, water grass with at least 1/4 inch of water after application, however stop watering before puddling occurs. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.
4. Water grass with at least 1/4 inch of water after application, however stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year.

Treat entire lawn; not just insect damaged areas. Do not allow traffic on lawn after application until granules are watered off lawn foliage and lawn has dried.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict according with directions giver. herewith.

SPREADER SETTING GUIDES \*\*

FOR AGWAY

Greenlawn Plus 21-5-7 Fertilizer With Diazinon Insecticide

Rate to deliver 3.6 lbs. per 1000 sq.ft.	Cyclone B1 Spinner Type	Ortho Spinner Type	Scotts Spinner Type	MTD Drop Type	Imperial Mark II Model 40 Drop Type	Scotts Drop Type
	Once Over	Once Over	Once Over	Once Over	Once Over	Once Over

\*\* NOTE: These settings are furnished as a guide only, age, condition of spreader, speed of operation and roughness of terrain can cause variation in rates applied.

CODE 86-