

## AGWAY GARDEN WEEDER II

### FOR HOME GARDEN USE ONLY

#### FOR USE IN ASPARAGUS & CANEBERRIES FOR THE CONTROL OF ANNUAL BROADLEAF & GRASSY WEEDS:

##### ACTIVE INGREDIENT:

Simazine (2-chloro-4, 6-bis (ethylamino) -s-triazine) . . . . . 4%

INERT INGREDIENTS: . . . . . 96%

TOTAL . . . . . 100%

### CAUTION

KEEP OUT OF REACH OF CHILDREN.

See additional precautionary statements.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE & DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

**ASPARAGUS** — Use only on established cutting beds. Apply 2 ounces (1/3 cup) per 100 sq. ft. in spring before weeds and asparagus emerge.

**BLUEBERRIES** — Make a single application of 1-5/6 — 3-2/3 oz (4.5 tablespoons to 9 tablespoons) per 100 sq. ft. in the spring before bud break and weed emergence. As an alternative procedure, apply 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the spring before bud break and weed emergence and follow with a second application of 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the fall. Do not apply when fruit is present.

To control quackgrass, apply 3-2/3 oz. (9 tablespoons) per 100 sq. ft. in the fall, or split the application by applying 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the fall and 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the early spring, when quackgrass is actively growing and before bud break.

**NOTE:** 1) On new plantings (less than 6 months old), use **one half** of the above rates; 2) Do not apply when fruit is present

**CANEERRIES (i.e. blackberries, boysenberries, loganberries, raspberries):** Make a single application of 1-5/6 — 3-2/3 oz (4.5 — 9 tablespoons) per 100 sq. ft. in the spring before bud break and weed emergence.

To control quackgrass, apply 3-2/3 oz. (9 tablespoons) per 100 sq. ft. in the fall, or split the application by applying 1-5/6 oz (4.5 tablespoons) per 100 sq. ft. in the fall and 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the spring when quackgrass is actively growing

**NOTE:** 1) On new plantings (less than 6 months old), use **one half** the above rates; 2) Do not apply when fruit is present.

**Established Nursery Stock** (See list below). Apply 1-5/6 — 2-3/4 oz (4.5 tablespoons to 6.75 tablespoons) per 100 sq. ft. in the fall or spring to nursery stock established at least one year after transplanting. Apply prior to weed emergence

Agway Garden Weeder II may be applied to these species of trees & shrubs, as directed above

AMERICAN ELM  
ARBORVITAE  
AUSTRIAN PINE  
BALSAM FIR  
BARBERRY  
BLUE SPRUCE  
BOXELDER  
BUSH HONEYSUCKLE  
CARAGANA  
COTONEASTER  
DOGWOOD

DOUGLAS FIR  
FRASER FIR  
HEMLOCK  
HONEY LOCUST  
JUNIPER  
MUGHO PINE  
NORWAY SPRUCE  
OREGON GRAPE  
(MAHONIA spp.)  
RED CEDAR  
RED OAK

RED PINE  
(NORWAY PINE)  
RED SPRUCE  
RUSSIAN OLIVE  
SCOTCH PINE  
SIBERIAN ELM  
WHITE CEDAR  
WHITE PINE  
WHITE SPRUCE  
YEW (TAXUS spp.)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin or clothing. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse

**First Aid:** In case of contact with eyes, flush with plenty of water for 15 minutes. Get medical attention if irritation persists. In case of contact with skin, wash with soap and water. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth, get medical attention.

### ENVIRONMENTAL HAZARDS

Do not contaminate domestic, livestock, or irrigation water supplies or lakes, streams or ponds. Do not contaminate water supplies by cleaning of equipment or disposal of wastes.


**DISCLAIMER:** Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

**NOTE:** This pesticide is to be sold **ONLY** in this original unbroken package.

EPA Reg. No. 8590-508

EPA Est. No. 8590-PA 1

AGWAY INC. — CHEMICAL DIVISION  
BOX 4933, SYRACUSE, NY 13221



34

**NET CONTENTS 1 LB.**