EDEM

FOR:

AGWAY GARDEN WEEDER II

DATE: 8/14/84

PCA 16 C"484C"1P1

RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND APPLICATION ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION.

NET CONTENTS
1 LB.

SURGHAY

SARDEN WEEDER II

For use in asparagus & caneberries for the control of annual broadleaf and grassy weeds:

ACTIVE INGREDIENT:

Simazine (2-chloro-4, 6-bis (ethylamino)

-s-triazine

4%

INERT INGREDIENTS:

96%

Total

100%

KEEP OUT OF REACH OF CHILDREN (12 pt)

CAUTION (18 pt)

See additional precautionary statements on side panel.

AGWAY INC. Chemical Division P. O. Box 4933

Syracuse, New York 13221

500

EPA Reg. No. 8590-508

EPA Est. No. 8590-PA-1

G'K.BY

•

SHEET NO. 1-5

PICA SECHARACIERS
ELITE - GECHARACIERS

ADVERTISING AND SALES PROMOTION SERVICES

GOPY

FOR:

AGWAY GARDEN WEEDER II

DATE: 8/14/84

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL (12 pt)

PROHIBITIONS: Do not contaminate or store near food, feed, fertilizers, seeds, insecticides, fungicides or other pesticides. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for use or storage in or around the home. Protect from excess heat.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION:

Since simazine enters weeds mainly through their roots, moisture is needed to move it into the root zone. This can be accomplished by rainfall or by watering in after application. Very dry soil conditions following application may necessitate shallow cultivation.

ASPARAGUS - Use only on established cutting beds. Apply 2 ounces (1/3 cup) per 100 sq. ft. in spring before weeds and asparagus emerge.

BLUEBERRIES - Make a single application of 1-5/6 - 3 2/3 :)
oz. (4.5 tablespoons to 9 tablespoons) per 100 sq. ft.
in the spring before bud break and weed emergence. As an alternative procedure, apply 1-5/6 oz. (4.5 tablespoons)
per 100 sq. ft. in the spring before bud break and weed emergence and follow with a second application of 1-5/6 oz.
(4.5 tablespoons) per 100 sq. ft. in the fall. Do not apply when fruit is present.

To control quackgrass, apply 3-2/3 oz (9 tablespoons) per 100 sq. ft. in the fall, or split the application by applying 1-5/6 oz (4.5 tablespoons) per 100 sq. ft. in the fall and 1-5/6 oz. (4.5 tablespoons) per 100 sq. ft. in the

, A K BA

SHEET NO. 2-5

COPY

- 94

FOR:

AGNAY GARDEN WEEDER II

DATE: 8/14/84

early spring, when quackgrass is actively growing and before bud break.

NOTE: 1) On new plantings (less than 6 months old), use one half of the above rates; 2) Do not apply when fruit is present.

CANEBERRIES (i.e. blackberries, boysenberries, loganberries, raspberries) - Make a single application of 1-5/6 - 3-2/3 oz (4.5 - 9 tablespoons) per 100 sq. ft. in the spring before bud break and weed emergence.

To control quackgrass, apply 3-2/3 oz (9 tablespoons) per 100 sq. ft. in the fall, or split the application by applying 1-5/6 oz (4.5 tablespoons) per 100 sq. ft. in the fall and 1-5/6 oz (4.5 tablespoons) per 100 sq. ft. in the spring when quackgrass is actively growing.

NOTE: 1) On new plantings (less than 6 months old), use one half the above rates; 2) Do not apply when fruit is present.

ESTABLISHED NURSERY STOCK (See list below) - Apply 1-5/6 - 2-3/4 cz (4.5 tablespoons to 6.75 tablespoons) per 100 sq. ft. in the fall or spring to nursery stock established at least one year after transplanting. Apply prior to weed emergence.

Agway Garden Weeder II may be applied to these species of trees and shrubs as directed above:

American Elm
Arborvitae
Austrian Pine
Balsam Fir
Barberry
Blue Spruce
Boxelder
Bush Honeysuckle
Caragana
Cotoneaster
Dogwood

Douglas Fir Fraser Fir Hemlock Honey Locust Juniper Mugho Pine Norway Spruce Oregon Grape (Mohonia spp) Red Cedar Red Oak

Red Pine
(Norway Pine)
Red Spruce
Russian Olive
Scotch Pine
Siberian Elm
White Cedar
White Pine
White Spruce
Yew (Taxus spp)

FICA 36 CHARACTERS ELITE - 48 CHARACTERS

TKBY

FOR: AGMAY GARDEN WEEDER II **DATE:** 8/14/84 PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin or clothing. Wash thoroughly after handling. Remove and wash contaminated clothing before reuse. First Aid: In case of contact with eyes, flush with plenty of water for 15 minutes. Get medical attention if irritation persists. In case of contact with skin, wash with soap and water. If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth, get medical attention. **ENVIRONMENTAL HAZARDS** Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Simazine is known to leach through soil, and has been found in ground water as a result of agricultural use. Users are advised not to apply in areas where soils are permeable: i.e., well drained, and which overlie shallow

SHEET NO. 4-5

POPY

DATE: 8/14/84 FOR: AGHAY GARDEN WEEDER II aquifers, particularly those currently being used for drinking water. Consult with the pesticide state lead agency for information regarding soil permeability and aguifer locations in your area. DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith. NOTE: This pesticide is to be sold ONLY in this original unbroken package. SHEET NO. 5-5